


Sampling Effort and Choice of Methods

Jacques H. C. Delabie, Brian L. Fisher,
Jonathan D. Majer, and Ian W. Wright



A number of studies have qualitatively assessed the efficacy of diverse methods for sampling rainforest ground-dwelling ants, including pitfall traps (e.g., Adis 1979), Winkler extraction (e.g., Olson 1991), baits (e.g., Fowler 1995), hand collection from quadrats (Room 1975), and “trap-nesting” using artificial nesting sites (Young 1986). However, very few studies have addressed the question of how many samples need to be taken in order to obtain a reasonably complete census of an ant community (e.g., Fisher 1999a). This deficiency of the literature results in part from the fact that the goal of most past ant sampling studies has been to provide a general inventory of the ant fauna of a region (e.g., Wilson 1959; Cover et al. 1990; Veerhagh 1990) rather than a rigorous measure of the underlying

biodiversity in terms of species numbers and relative abundances.

The qualitative approach may well be adequate when the aim is merely to provide a list of the species present. However, if ants are to be used as bioindicators of some aspect of the environment, or if a rigorous census of species is desired, then richness and abundance measures must be described explicitly per unit area or per unit of sampling effort. In such cases, it is necessary to know whether the samples are or are not capturing a reasonably high proportion of the ant species present; if they are not, then it is necessary to estimate what proportion of the total ant fauna is being sampled.

A single sampling method is unlikely to capture all the ants present in the litter or in other

parts of the ground stratum. To overcome this problem, researchers often use a combination of complementary sampling procedures. Combinations that have been used to sample rainforest ground-dwelling ants include Winkler extraction of both litter and soil (Belshaw and Bolton 1994a); Winkler extraction plus pitfall traps (Olson 1991); Winkler extraction plus pitfall traps plus hand collecting (Andersen and Majer 1991; Fisher 1996a, 1998, 1999b); pitfall traps plus hand collecting (Jackson 1984); pitfall traps plus baits (Fowler 1995); and Berlese funnels plus baits (Levings 1983). In this chapter we use data from comprehensive studies of the ant communities in Brazilian cocoa plantations to address (1) the optimal combination of sampling methods for maximizing the species count of rainforest ground-dwelling ants; and (2) the relationship between the size of individual litter samples and the number of species obtained by Berlese funnel or Winkler extraction methods.

Methods

Field work was carried out on the grounds of the Centre for Cocoa Research (CEPLAC), Ilheus, Bahia ($14^{\circ}45'S$, $39^{\circ}13'W$), Brazil. The CEPLAC site formerly consisted of primary Atlantic rainforest (Mata Atlantica), although most of it is now planted with cocoa. Cocoa plantations provide a habitat that retains many features of the original rainforest, and the ground-dwelling ant fauna retains a high degree of similarity to the fauna of the original habitat (Belshaw and Bolton 1993; Young 1986; Delabie et al. 2000).

Sampling Methods Experiment

A 1-ha plantation of regularly planted 20-year-old cocoa trees, shaded with *Erythrina fusca*, was divided into three rows of six cells, each measuring $23.5\text{ m} \times 23.5\text{ m}$. For each sampling method, three sample points were randomly selected in each cell, resulting in a total of 54

samples for each method for the entire plot. Sampling was originally designed to census the ant fauna from the soil, the litter, the ground surface, the tree trunks, and the canopy; the preliminary results have been reported in Delabie et al. (1994). Here we consider only those sampling procedures that are relevant to the soil and litter stratum. The 17 sampling methods used were as follows:

1. Small soil samples. Cubes of soil measuring 15 cm on a side were dug up, broken open, sieved, and then inspected on a white surface so that ants could be manually removed.
2. Large soil samples. As in (1), except that the sides of the cubes were 30 cm across.
3. Berlese funnel samples. Samples of litter measuring 1 m^2 were collected and placed in a funnel for 24 hours.
4. Winkler extraction samples. As in (3), except that the litter was sieved and then placed in Winkler sacks for 24 hours.
5. Pitfall traps (24-hour). A 75-mm-internal-diameter pitfall trap, containing water and a drop of detergent, was placed out for 24 hours.
6. Pitfall traps (7-day). As in (5), except that the traps contained a mixture of ethanol and glycerol and were placed out for 7 days.
7. Sardine bait (4-hour). Small pieces of sardine were placed on a square of tissue paper, and the ants that were attracted were collected after 4 hours.
8. Sardine bait (24-hour). As in (7), except that the baits were inspected after 24 hours.
9. Meat bait (24-hour). Small pieces of meat were placed on a square of tissue paper, and the ants that were attracted were collected after 24 hours.
10. Cassava flour bait (4-hour). A small pile of coarse cassava flour was placed on a

- square of tissue paper, and the ants that were attracted were collected after 4 hours.
11. Cassava flour bait (24-hour). As in (10), except that the baits were inspected after 24 hours.
 12. Sugar bait (4-hour). A small amount of dilute sugar solution was placed on a square of tissue paper, and the ants that were attracted were collected after 4 hours.
 13. Sugar bait (24-hour). As in (12), except that the baits were inspected after 24 hours.
 14. Orange peel bait (4-hour). A small piece of orange peel was placed on a square of tissue paper, and the ants that were attracted were collected after 4 hours.
 15. Orange peel bait (24-hour). As in (14), except that the baits were inspected after 24 hours.
 16. Dead wood inspection. A rotting branch near the sampling point was cut open and the ants within were collected.
 17. Dried cocoa pod inspection. A rotting cocoa pod near the sampling point was cut open and the ants within were collected.

A number of additional sampling methods were used to collect ants from other strata in the plantation. These included chemical knock-down and beating of trees, meat and sardine baiting in trees, inspection of dried cocoa pods on trees, inspection of fallen epiphytic bromeliads, and direct observation over fixed time intervals. The ant collections resulting from these methods are outside the scope of this chapter and will be reported in a subsequent publication.

Extended Sampling Experiment

To investigate the effect of number of samples on the number of species obtained, an extended set of soil and litter samples were collected in a nearby plantation of 60-year-old cocoa. This area differs from the 20-year-old plantation described previously in two important ways:

(1) the cocoa is grown in a less “manicured” way, is shaded by around 15 species of native overstory rainforest trees, and thus constitutes a more structurally diverse habitat than the 20-year-old plantation; and (2) no pesticides have been applied to it for 30 years. In this plantation, 500 Winkler extractions were collected from 1-m² samples of litter taken from randomized points in a 0.87-ha area.

Sample Size Experiment, 60-Year-Old Brazilian Cocoa Plantation

To investigate the influence of litter sample size on the number of species obtained, a set of samples was taken from the same 60-year-old plantation described previously, but using different litter sample sizes. Samples of 0.01 m², 0.04 m², 0.25 m², and 1 m², each replicated 20 times, were collected and extracted by Berlese funnels over 24 hours. An identical sampling regimen using Winkler extraction methods was also undertaken, except that additional samples of 2 m² were also taken.

Data Analysis

All ants from each experiment were initially sorted to the level of morphospecies, then identified to genus and, where possible, to species. For each sampling method, we constructed a matrix of individual ant species by individual samples and determined the frequency of each species in the samples. Standard diversity analyses were then carried out on the resulting data sets. See Chapter 13 and Magurran (1988) for additional background on the statistical analyses described in this section.

The incidence-based coverage estimator (ICE) (Lee and Chao 1994; Chazdon et al. 1998) and the first-order jackknife estimator (Heltshe and Forrester 1983) are nonparametric approaches for estimating species richness in the local community from which the samples were taken, that is, for estimating how many

species (including those not collected) are present in the sampled community. Calculation of the ICE is based on the number of species found in ten or fewer sampling units, whereas calculation of the jackknife is based on the observed frequency of unique species. To estimate what proportion of the total species richness was collected by each of the methods employed in the current study, plots of cumulative species-per-sample curves were generated in which species accumulation was plotted as a function of the number of samples taken. Three values were plotted for each succeeding sample: the observed number of species, the ICE of the total number of species present, and the first-order jackknife estimate of the total number of species present. For species-accumulation curves, sample order was randomized 100 times and the means of the ICE and jackknife estimates were computed for each succeeding sample station using the program EstimateS (Colwell 1997; see also Colwell and Coddington 1994; Chazdon et al. 1998).

The asymptote of a species-accumulation curve (i.e., the value that the curve approaches as a limit) is interpreted as the total number of species present at the sample site, including those that were not collected. In the current study, the asymptote of the observed species-accumulation curve was calculated with EstimateS (Colwell 1997) using the two-parameter Michaelis-Menten (M-M) equation (Colwell and Coddington 1994) and the maximum likelihood method of Raaijmakers (1987), which is based on the Eadie-Hofstee transformation of the M-M equation. The observed number of ant species and the proportion of the M-M asymptote represented by this number were evaluated for different sample sizes for each of the sampling methods.

The numbers of species collected by different combinations of the various sampling methods have important implications for future sampling studies. To elucidate the optimal combination of

sampling methods tested in the sampling methods experiment, first a table was prepared listing the total number of species sampled by each method. The method that sampled the most species was identified, and combinations including one and two additional sampling procedures were assessed in order to elucidate the optimal combinations of two and three sampling methods for maximizing the species count.

To evaluate the influence of the size of the litter sample from which ants were collected by Berlese funnel or Winkler extraction, the mean number of individual ants and the number of species per extraction were calculated; the total number of species from 20 samples of a particular area was also calculated.

Results and Discussion

Interpretation of species richness estimates should take into account a number of factors. A species-accumulation curve is specific to the area of the survey, the season or year, and the collecting techniques employed. The use of additional collecting methods, or a survey in a different area or season at the same site, would most certainly collect additional species. The actual number of species in an area at a given time is of course finite, but, in most cases, exhaustive sampling is not physically or logistically possible. If an observed or estimated species-accumulation curve indicates a decrease in the rate of species accumulation across the number of samples collected, then, for the particular methods employed, that number of samples is arguably adequate for estimating the species richness in the area or transect surveyed. Conversely, if the curve continues to rise rapidly for the number of samples collected, then more intensive sampling may be necessary to obtain an adequate measure of the diversity at that site.

The number of samples sufficient for achieving a high level of species completeness is thus

practically defined as the point at which the accumulation curve shows an adequate decrease in species accrual. The principal problems with this “sufficient-sampling” definition are the lack of a single asymptote for diverse taxa and the difficulty in quantifying “an adequate decrease in species accrual.” One practical solution to the latter problem is to sample until a certain percentage—say 80%—of the estimated species are obtained, based on ICE and jackknife estimates of the total number of species that occur in the plot or transect. An alternative approach is to continue to sample until additional sampling effort is predicted to achieve a negligible increase in the number of species sampled. In this approach, the increase in species obtained with additional sampling must be weighed against both the cost of sorting and identifying additional specimens and the relative importance of identifying the full complement of the species at the site. Under either approach, species-accumulation curves can be calculated by randomizing sample accumulation order and using asymptotic or nonasymptotic functions (Colwell and Coddington 1994), and the predicted species richness values can be extrapolated using the ICE, jackknife, and M-M estimation techniques.

Sampling Methods Experiment

The 17 sampling methods yielded a total of 134 ant species. In the total of 918 samples (17 methods \times 54 samples per method), most of these ground-dwelling ant species are extremely rare, with 43 species collected only once, 18 species collected twice, and 34 species collected in up to 10 samples; only 39 species occurred in 11 or more samples.

The number of species sampled by the 17 different methods is shown in Table 10.1. Winkler extraction sampled the greatest number of species (63), followed by Berlese funnels (48), inspection of dead wood (45), small soil samples (42), and pitfall traps. Pitfall traps run for 7

Table 10.1 Actual Number of Ant Species Sampled by the 17 Methods Described in the Text

Sampling Method	Number of Species	Rank ^a
Winkler extraction samples	63	1
Berlese funnel samples	48	2
Dead wood inspection	45	3
Small soil samples	42	4
Pitfall traps (7-day)	40	5
Pitfall traps (24-hour)	27	6
Large soil samples	26	7
Sardine bait (24-hour)	20	8.5
Orange peel bait (24-hour)	20	8.5
Sardine bait (4-hour)	19	10.5
Orange peel bait (4-hour)	19	10.5
Sugar bait (4-hour)	18	12
Dried cocoa pod inspection	17	13
Cassava flour bait (4-hour)	16	14
Meat bait (24-hour)	15	15
Cassava flour bait (24-hour)	14	16
Sugar bait (24-hour)	11	17

^aMethods are ranked from 1 (most species sampled) through 17 (fewest species sampled).

days yielded more ant species than those run for 24 hours (40 versus 27 species). Large soil samples yielded only 26 species and dead cocoa pods only 17 species, while the species counts from the baiting methods never exceeded 20. Baiting for longer periods of time did not necessarily yield more species; indeed, with the cassava and the sugar baits, a lower number of species was obtained after the longest baiting period.

The species-accumulation curve plots are shown for the individual methods in Figs. 10.1a–q, and for all methods combined in Fig. 10.1r. These graphs also show jackknife and ICE estimators of species richness based on successively larger numbers of samples. The agreement between the predicted M-M asymptotic values and the actual maximum value encountered in the 54 samples is shown in Table 10.2. This table




Figure 10.1. Assessment of each of 17 leaf litter ant sampling methods in Brazil. The lower (thick) species-accumulation curve plots the observed number of species as a function of the number of stations sampled. The upper curves display the nonparametric first-order jackknife estimator (dashed) and ICE (solid). The estimated total species richness is based on successively larger numbers of samples from the data set (Heltsh and Forrester 1983; Lee and Chao 1994). Curves are plotted from the means of 100 randomizations of sample accumulation order.

Table 10.2 Observed Number of Ant Species Evaluated at Different Sample Sizes for Each of the 17 Sampling Methods^a

Methods	Observed Species Richness after:					Estimated Species Richness ^b		
	10 Samples	20 Samples	30 Samples	40 Samples	All (54) Samples	ICE	Jack-knife	M-M
Small soil samples	14.0 (20.9)	22.4 (33.5)	25.3 (37.9)	29.6 (44.3)	42 (62.8)	88.5	64.6	66.9
Large soil samples	11.9 (39.1)	16.4 (53.9)	20.0 (65.8)	22.7 (74.5)	26 (85.5)	49.0	37.8	30.4
Berlese funnel samples	21.8 (39.4)	31.2 (56.4)	37.8 (68.4)	42.4 (76.7)	48 (86.8)	65.7	64.7	55.3
Winkler extraction samples	29.3 (42.2)	41.1 (59.1)	49.1 (70.6)	55.0 (79.1)	63 (90.6)	108.1	90.5	69.5
Pitfall traps (24-hour)	12.2 (35.9)	17.4 (51.1)	20.8 (61.1)	23.9 (70.2)	27 (79.4)	40.0	37.8	34.0
Pitfall traps (7-day)	17.3 (36.5)	24.5 (51.7)	30.0 (63.2)	34.4 (72.4)	40 (84.3)	72.8	59.6	47.5
Sardine bait (4-hour)	7.3 (34.6)	10.5 (49.4)	13.2 (62.2)	15.8 (74.6)	19 (89.8)	68.5	30.8	21.2
Sardine bait (24-hour)	9.1 (37.3)	13.0 (53.4)	15.8 (64.7)	17.8 (73.2)	20 (82.1)	29.8	27.9	24.4
Cassava flour bait (24-hour)	6.1 (36.4)	8.9 (52.8)	10.8 (64.1)	12.2 (72.7)	14 (83.2)	23.8	19.9	16.8
Cassava flour bait (4-hour)	6.2 (27.5)	9.3 (41.8)	11.8 (52.8)	13.9 (62.2)	16 (71.6)	24.6	22.9	22.4
Meat bait (24-hour)	6.5 (35.0)	9.7 (52.4)	11.8 (64.2)	13.4 (72.6)	15 (81.3)	18.3	19.9	18.4
Sugar bait (4-hour)	8.7 (41.0)	12.0 (56.7)	14.3 (68.0)	16.0 (75.9)	18 (85.4)	27.0	24.9	21.1
Sugar bait (24-hour)	5.2 (39.3)	7.3 (54.9)	8.9 (66.9)	10.0 (75.5)	11 (83.0)	12.7	13.9	13.3
Orange peel bait (4-hour)	7.1 (31.0)	10.8 (47.1)	13.8 (60.4)	16.1 (70.0)	19 (82.9)	34.9	28.8	22.9
Orange peel bait (24-hour)	8.1 (31.8)	12.0 (47.2)	14.8 (58.1)	17.2 (67.5)	20 (78.5)	38.2	30.8	25.5
Dead wood inspection	17.6 (27.9)	27.6 (43.6)	34.1 (54.0)	39.2 (62.0)	45 (71.3)	71.0	65.6	63.1
Dried cocoa pod inspection	7.1 (32.7)	11.2 (51.2)	13.6 (62.4)	15.2 (69.8)	17 (77.9)	22.8	22.9	21.8
All methods	75.0 (55.0)	95.8 (70.3)	110.1 (80.7)	121.5 (89.1)	134 (98.3)	188.9	176.2	136.4

^aNumber of species represents the mean of 100 randomizations of sample pooling order.

^bICE, incidence-based coverage estimator; jackknife, first-order jackknife estimator; M-M, Michaelis-Menten asymptote (the percentage of the M-M asymptote is given in parentheses in the first five columns).

also shows the ICE and jackknife asymptote values, as well as the predicted proportions (as percentages) of the M-M asymptotic value that would be obtained if 10, 20, 30, 40, and all 54 samples were taken.

Depending on the sampling method, the percentage of the M-M asymptote value that was obtained varied from 20.9 to 42.2, 33.5 to 59.1, 44.3 to 70.6, 37.9 to 79.1, and 62.8 to 90.6 when 10, 20, 30, 40, and 54 samples were collected, respectively. In all cases, Winkler extraction obtained the highest percentage of the asymptote value and small soil samples the lowest.

There is reasonable agreement between the percentages of the asymptotes obtained for the various sampling methods, which were on average 34.6, 50.4, 61.8, 69.8, and 81.0 for 10, 20, 30, 40, and 54 samples, respectively.

For all methods combined, 70% of the asymptote value was collected by 20 samples, whereas for each individual method, a much lower percentage (33.5–59.1%) was collected by 20 samples, with some methods obviously performing much better than others. Thus, in the context of the total survey, increasing the number of samples from each individual

method did not necessarily add new species to the total census, although for particular methods, species that were rarely collected by one method were common in another.

Considering that the study area was sampled by 17 different methods and that a relatively large number of samples was taken, the total species count probably represents a nearly complete census. This conclusion is additionally supported by an asymptote value of 136.4 species (Fig. 10.1r) for the curve for all methods combined, which implies that the combination of 54 samples from all 17 methods obtained 98.3 percent of the asymptote.

Extended Sampling Experiment

The species-accumulation curves for the 500 Winkler extraction samples are shown in Fig. 10.2. As above, this graph also shows ICE, jackknife, and M-M estimators of species richness based on successively larger numbers of samples. The agreement between the predicted M-M asymptotic value and the actual maximum value encountered in the 500 samples is shown in Table 10.3. This table also shows the ICE and jackknife asymptotes, as well as the predicted percentages of the M-M asymptotic value that would be obtained if 10, 20, 30, 40, 100, 200, 300, 400, and 500 samples were taken.

If we were to compare the M-M asymptote value of 136.4 species for this extended sampling experiment (carried out in the 60-year-old cocoa plantation) with the asymptote value of 69.5 species obtained for only the Winkler extraction samples from the sampling methods experiment (carried out in the 20-year-old cocoa plantation), we might be led to conclude that ant species richness is higher in the 60-year-old cocoa plantation. In fact, this disparity is probably an artifact of the greater number of Winkler extraction samples that were taken in the extended sampling experiment, in which an average of 67.7 species was collected with 54 pooled samples. This value is quite close to the




Figure 10.2. Assessment of Winkler sampling methods in Brazil. Lines follow the same convention as in Fig. 10.1.

observed total of 63 species for 54 Winkler sack samples in the sampling methods experiment. When the species-accumulation curve for the 54 Winkler sacks in the sampling methods experiment is extrapolated out to 500 samples using the logarithmic equation of Soberón and Llorente (1993), 109.9 species are predicted. This is also quite close to the 107 species observed in the extended sampling experiment. But perhaps the most telling argument against a significant difference in species richness between the 20- and 60-year-old plantations is the similarity in richness estimates of 134.6 species for the former (based on all sampling methods combined) and 136.4 for the latter (based on the full complement of 500 Winkler extraction samples). It is particularly noteworthy that the first 100 samples yielded over three-quarters of the species that were ultimately sampled by this procedure, and that the last 200 samples yielded very few additional species.

Sample Size Experiment

For both the Berlese funnel and Winkler sack samples, number of individual ants, total number of ant species, and mean number of ant species per sample generally increased with increasing litter sample size. However, although

Table 10.3 Observed Number of Ant Species Evaluated at Different Sample Sizes for the Extended Winkler Sampling Experiment^a

Observed Species Richness after Following Number of Samples:										Estimated Species Richness ^b		
10	20	30	40	100	200	300	400	All (500)	ICE	Jackknife	M-M	
36.5 (35.2)	50.0 (48.2)	56.3 (54.3)	62.1 (60.0)	79.5 (76.7)	92.7 (89.5)	100.1 (96.6)	104.5 (100.8)	107 (103.3)	112.0	117.0	103.6	

^aNumber of species represents the mean of 100 randomizations of sample pooling order.

^bICE, incidence-based coverage estimator; jackknife, first-order jackknife estimator; M-M = Michaelis-Menten asymptote (the percentage of the M-M asymptote is given in parentheses in the first nine columns).

the number of individuals increased quite dramatically with increasing sample size, the increase in mean number of species per sample was far less pronounced. Increasing the sample size from 0.25 m² to 1 m², and certainly from 1 m² to 2 m², is associated with a very limited return for a costly additional investment in effort.

Based on the results of the sample size experiment, when sufficient sampling devices are available it is probably generally more efficient to take a greater number of smaller samples than to take a lesser number of larger samples. Since a single Winkler sack can generally hold the sieved litter from a 1-m² sample, this is probably the most appropriate sample size for most situations.

Complementarity of Sampling Methods

Table 10.4 lists the combinations of two and three sampling methods that produced the largest numbers of ant species in the sampling methods experiment. In each case, only combinations that produced the four highest species counts are shown. Winkler extraction is an element in all combinations, along with inspection of dead wood, small soil samples, pitfall traps (7-day), and Berlese funnels. The various two-

method combinations captured species totals ranging from 59 to 65% of the M-M asymptote calculated for all methods combined (Table 10.2). Species totals for three-method combinations ranged from 73 to 77% of the M-M

Table 10.4 Combinations of Two and Three Sampling Methods That Obtained the Maximum Number of Ant Species in the Sampling Methods Experiment

Combination of Sampling Methods	N ^a
Winkler sack samples + small soil samples + inspection of dead wood	105
Winkler sack samples + inspection of dead wood + pitfall traps (7-day)	104
Winkler sack samples + pitfall traps (7-day) + small soil samples	103
Winkler sack samples + small soil samples + Berlese funnel samples	99
Winkler sack samples + inspection of dead wood	88
Winkler sack samples + small soil samples	87
Winkler sack samples + pitfall traps (7-day)	84
Winkler sack samples + Berlese funnel samples	80
Total number of litter species sampled by all methods	134
Total number of species from all strata	167


^aN, number of species.

asymptote calculated for all methods combined. A similar exercise (results not shown) was performed with combinations of four sampling methods. Generally speaking, the combinations that maximized species counts are simply permutations of the methods listed in Table 10.4. The combination of methods that produced the maximum species count (117 species) was Winkler sack samples + small soil samples + inspection of dead wood + pitfall traps (7-day). This value of 117 species is 86% of the asymptote value calculated for all methods combined (Table 10.2).

Conclusion

The choices of what collecting methods to use and how many samples to collect are dependent

on the intended species completeness of the proposed inventory, that is, on what proportion of the ant fauna the inventory intends to survey. The results presented here demonstrate that Winkler extraction is the most efficient method for surveying leaf litter ants and therefore that this method should be included in all ground-dwelling ant inventory protocols. If a second method is also to be used, we recommend pitfall traps. The total number of samples to collect should be determined both by the alpha diversity of the inventory site and by the level of species completeness that is necessary to achieve the project's goals, but, based on the results reported here, in most situations we recommend taking 20 1-m² Winkler samples for areas comparable in size to the 1-ha 20-year-old cocoa plantation.



Literature Cited

- Abensperg-Traun, M., and D. Steven. 1995. The effects of pitfall trap diameter on ant species richness (Hymenoptera: Formicidae) and species composition of the catch in a semi-arid eucalypt woodland. *Australian Journal of Ecology* 20:282–287.
- _____. 1997. Ant- and termite-feeding in Australian mammals and lizards: A comparison. *Australian Journal of Ecology* 22:9–17.
- Abensperg-Traun, M., G. W. Arnold, D. E. Steven, G. T. Smith, L. Atkins, J. J. Viveen, and M. Gutter. 1996. Biodiversity indicators in semi-arid, agricultural Western Australia. *Pacific Conservation Biology* 2:375–389.
- Adams, E. S. 1994. Territory defense by the ant *Azteca trigona*: Maintenance of an arboreal ant mosaic. *Oecologia* 97:202–208.
- Adams, E., and W. Tschinkel. 1995. Density-dependent competition in fire ants: Effects on colony survivorship and size variation. *Journal of Animal Ecology* 64:315–324.
- Adis, J. 1979. Problems of interpreting arthropod sampling with pitfall traps. *Zoologisher Anzeiger* 202:177–184.
- Adis J., Y. D. Lubin, and G. G. Montgomery. 1984. Arthropods from the canopy of inundated and terra firme forests near Manaus, Brazil, with critical considerations on the pyrethrum-fogging technique. *Studies on Neotropical Fauna and Environment* 4:223–236.
- Adis, J., J. W. de Moraes, and H. Guimarães de Mesquita. 1987. Vertical distribution and abundance of arthropods in the soil of a Neotropical secondary forest during the rainy season. *Studies on Neotropical Fauna and Environment* 22:189–197.
- Agosti, D. 1990. Review and reclassification of *Cataglyphis* (Hymenoptera, Formicidae). *Journal of Natural History* 24:1457–1505.
- _____. 1991. Revision of the oriental ant genus *Cladomyrma*, with an outline of the higher classification of the Formicinae (Hymenoptera:

- Formicidae). *Systematic Entomology* 16:293–310.
- _____. 1992. Revision of the ant genus *Myrmoteras* of the Malay Archipelago (Hymenoptera, Formicidae). *Revue Suisse de Zoologie* 99:405–429.
- _____. 1994a. The phylogeny of the ant tribe Formicinae (Hymenoptera: Formicidae) with the description of a new genus. *Systematic Entomology* 19:93–117.
- _____. 1994b. A revision of the South American species of the ant genus *Probolomyrmex* (Hymenoptera: Formicidae). *Journal of the New York Entomological Society* 102:429–434.
- Agosti, D., J. Majer, L. Alonso, and T. R. Schultz (eds.). 2000. *Sampling Ground-dwelling Ants: Case Studies from the Worlds' Rain Forests*. Curtin University of Technology, Perth, Western Australia.
- Agosti, D., M. Maryati, and C. Y. C. Arthur. 1994. Has the diversity of tropical ant fauna been underestimated? An indication from leaf litter studies in a West Malaysian lowland rain forest. *Tropical Biodiversity* 2:270–275.
- Agosti, D., J. Moog, and U. Maschwitz. 1999. Revision of the Oriental plant-ant genus *Cladomyrma*. *American Museum Novitates* 3283, December 8, 1999.
- Alexander, R. D. 1974. The evolution of social behavior. *Annual Review of Ecology and Systematics* 5:325–383.
- Allen, C., R. Lutz, and S. Demarais. 1995. Red imported fire ant impacts on northern bobwhite populations. *Ecological Applications* 5:632–638.
- Allen, G. E., and W. F. Buren. 1974. Microsporidian and fungal diseases of *Solenopsis invicta* Buren in Brazil. *Journal of the New York Entomological Society* 82:125–130.
- Allen, G. E., and A. Silveira-Guido. 1974. Occurrence of microsporidia in *Solenopsis richteri* and *Solenopsis* sp. in Uruguay and Argentina. *Florida Entomologist* 57:327–329.
- Allen, M. F., J. A. MacMahon, and D. C. Andersen. 1984. Reestablishment of Endogonaceae on Mount St. Helens: Survival of residuals. *Mycologia* 76(6):1031–1038.
- Andersen, A. N. 1983. A brief survey of ants of Glenaladale National Park, with particular reference to seed-harvesting. *Victorian Naturalist* 100:233–237.
- _____. 1986a. Diversity, seasonality and community organization of ants at adjacent heath and woodland sites in southeastern Australia. *Australian Journal of Zoology* 34:53–64.
- _____. 1986b. Patterns of ant community organization in mesic southeastern Australia. *Australian Journal of Ecology* 11:87–99.
- _____. 1988. Immediate and longer-term effects of fire on seed predation by ants in sclerophyllous vegetation of southeastern Australia. *Australian Journal of Ecology* 13:285–293.
- _____. 1990. The use of ant communities to evaluate change in Australian terrestrial ecosystems: A review and a recipe. *Proceedings of the Ecological Society of Australia* 16:347–357.
- _____. 1991a. Parallels between ants and plants: Implications for community ecology. Pp. 539–558. In C. R. Huxley and D. F. Cutler (eds.), *Ant-Plant Interactions*. Oxford University Press, Oxford.
- _____. 1991b. Seed-harvesting by ants in Australia. Pp. 493–503. In C. R. Huxley and D. F. Cutler (eds.), *Ant-Plant Interactions*. Oxford University Press, Oxford.
- _____. 1991c. Sampling communities of ground-foraging ants: Pitfall catches compared with quadrat counts in an Australian tropical savanna. *Australian Journal of Ecology* 16:273–279.
- Andersen, A. N. 1991d. Responses of ground-foraging ant communities to three experimental fire regimes in a savanna forest of tropical Australia. *Biotropica* 23:575–585.
- _____. 1992. The rainforest ant fauna of the northern Kimberley region of Western Australia (Hymenoptera: Formicidae). *Journal of the Australian Entomological Society* 31:187–192.
- _____. 1995. A classification of Australian ant communities, based on functional groups which parallel plant life-forms in relation to stress and disturbance. *Journal of Biogeography* 20:15–29.
- _____. 1996. Fire ecology and management. Pp. 179–195. In C. M. Finlayson and I. Von Oertzen (eds.), *Landscape and Vegetation Ecology of the Kakadu Region, Northern Australia*. Kluwer Academic Publishers, Amsterdam.
- _____. 1997a. Functional groups and patterns of organization in North American ant communities: A comparison with Australia. *Journal of Biogeography* 24:433–460.
- _____. 1997b. Using ants as bioindicators: Multi-scale issues in ant community ecology. *Conservation Ecology* [online] 1, Article 8.
- Andersen, A. N., and R. E. Clay. 1996. The ant fauna of Danggali Conservation park in semi-arid

- South Australia: A comparison with Wyperfield (Vic.) and Cape Arid (W.A.) National Parks. *Australian Journal of Entomology* 35:289–295.
- Andersen, A. N., and J. D. Majer. 1991. The structure and biogeography of rainforest ant communities in the Kimberley region of northwestern Australia. Pp. 333–346. In N. L. McKenzie, R. B. Johnston, and P. J. Kendrick (eds.), *Kimberley Rainforests of Australia*. Surrey Beatty and Sons, Chipping Norton, NSW.
- Andersen, A. N., and M. E. McKaige. 1987. Ant communities at Rotamah Island, Victoria, with particular reference to disturbance and *Rhytidoponera tasmaniensis*. *Proceedings of the Royal Society of Victoria* 99:141–146.
- Andersen, A. N., and A. D. Patel. 1994. Meat ants as dominant members of Australian ant communities: An experimental test of their influence on the foraging success and forager abundance of other species. *Oecologia* 98:15–24.
- Andersen, A. N., and H. Reichel. 1994. The ant (Hymenoptera: Formicidae) fauna of Holmes Jungle, a rainforest patch in the seasonal tropics of Australia's Northern Territory. *Journal of the Australian Entomological Society* 33:153–158.
- Andersen, A. N., and A. V. Spain. 1996. The ant fauna of the Bowen Basin, in the semi-arid tropics of central Queensland (Hymenoptera: Formicidae). *Australian Journal of Entomology* 35:213–221.
- Andersen, A. N., and A. L. Yen. 1985. Immediate effects of fire on ants in the semi-arid mallee region of north-western Victoria. *Australian Journal of Ecology* 10:25–30.
- . 1992. Canopy ant communities in the semi-arid Mallee region of North-western Victoria. *Australian Journal of Zoology* 40:205–214.
- Andersen, A. N., M. S. Blum, and T. M. Jones. 1991. Venom alkaloids in *Monomorium "rothsteini"* Forel repel other ants: Is this the secret to success by *Monomorium* in Australian ant communities? *Oecologia* 88:157–160.
- Andersen, A. N., S. Morrison, and L. Belbin. 1996. The Role of Ants in Minesite Restoration in the Kakadu Region of Australia's Northern Territory, with Particular Reference to Their Use as Bio-indicators. Final Report to the Environmental Research Institute of the Supervising Scientist, Australia.
- Andrade, M. L. de. 1998. Fossil and extant species of *Cylindromyrmex* (Hymenoptera: Formicidae). *Revue Suisse de Zoologie* 105(3):581–664.
- Andrade, M. L. de and C. Baroni Urbani. 1999. Diversity and adaption in the ant genus *Cephalotes*, past and present (Hymenoptera, Formicidae). *Stuttgarter Beiträge zur Naturkunde, Serie B* 271:1–889.
- Andrewartha, H., and L. Birch. 1954. The distribution and abundance of animals. University of Chicago Press, Chicago.
- Arnett, R. H., Jr. 1985. *American Insects*. Van Nostrand Reinhold, New York.
- Arnett, R. H., Jr., and M. E. Arnett. 1990. *The Naturalists Directory and Almanac (International): An Index to Contemporary Naturalists of the World and Their Special Interests*, 45th ed. Sandhill Crane Press, Gainesville, Florida.
- Arnett, R. H., Jr., and G. A. Samuelson. 1986. *The Insect and Spider Collections of the World*. E. J. Brill/Flora and Fauna Publications, Gainesville, Florida.
- Arnett, R. H., Jr., G. A. Samuelson, and G. M. Nishida. 1993. *The Insect and Spider Collections of the World*, 2nd ed. Sandhill Crane Press, Gainesville, Florida.
- Arnoldi, K. V. 1930. Studien über die Systematik der Ameisen. IV. *Aulacopone*, eine neue Ponerinengattung (Formicidae) in Russland. *Zoologischer Anzeiger* 89:139–144.
- . 1970. Review of the ant genus *Myrmica* (Hymenoptera, Formicidae) in the European part of the USSR. *Zoologicheskii Zhurnal* 49:1829–1844. [In Russian.]
- . 1975. A review of the species of the genus *Stenamma* (Hymenoptera, Formicidae) of the USSR and description of new species. *Zoologicheskii Zhurnal* 54:1819–1829. [In Russian.]
- . 1976a. Review of the genus *Aphaenogaster* (Hymenoptera, Formicidae) in the USSR. *Zoologicheskii Zhurnal* 55:1019–1026. [In Russian.]
- . 1976b. Ants of the genus *Myrmica* Latr. from Central Asia and the southern Kazakhstan. *Zoologicheskii Zhurnal* 55:547–558. [In Russian.]
- . 1977. Review of the harvester ants of the genus *Messor* (Hymenoptera, Formicidae) in the fauna of the USSR. *Zoologicheskii Zhurnal* 56:1637–1648. [In Russian.]
- Atsatt, P. R. 1981. Lycaenid butterflies and ants: Selection for enemy-free space. *American Naturalist* 118:538–654.
- Autuori, M. 1942. Contribuição para o conhecimento da saúva (*Atta* spp. Hymenoptera: Formicidae).

- III. Excavação de um saúveiro (*Atta sexdens rubropilosa* Forel, 1908). *Archivos do Instituto de Biolóico*, São Paulo 13:137–148.
- Bailey, I. W. 1920. Some relations between ants and fungi. *Ecology* 1:174–189.
- _____. 1922a. Notes on neotropical ant-plants. I. *Cecropia angulata*, sp. nov. *Botanical Gazette* 74:369–391.
- _____. 1922b. The anatomy of certain plants from the Belgian Congo, with special reference to Myrmecophytism. *Bulletin of the American Museum of Natural History* 45:585–622, plates 26–45.
- Balazy, S., A. Lenoir, and J. Wisniewski. 1986. *Aegeritella rousillonensis* n. sp. (Hymenomycetales, Blastosporae), une espèce nouvelle de champignon épizoïque sur les fourmis *Cataglyphis cursor* (Fonscolombe) (Hymenoptera, Formicidae) en France. *Cryptogamie, Mycologie* 7:37–45.
- Banschbach, V. S., and J. M. Herbers. 1996. Complex colony structure in social insects. I. Ecological determinants and genetic consequences. *Evolution* 50:285–297.
- Baroni Urbani, C. 1968. Über die eigenartige Morphologie der männlichen Genitalien des Genus *Diplorhoptrum* Mayr und die taxonomischen Schlussfolgerungen. *Zeitschrift für Morphologie der Tiere* 63:63–74.
- Baroni Urbani, C. 1969. Gli *Strongylognathus* del gruppo *huberi* nell'Europa occidentale: Saggio di una revisione basata sulla casta operaia (Hymenoptera Formicidae). *Bulletino de la Società Entomologica Italiana* 99–101:132–168.
- _____. 1975a. Primi reperti del genere *Calyptomyrmex* Emery nel subcontinente Indiano. *Entomologica Basiliensis* 1:395–411.
- _____. 1975b. Contributo alla conoscenza dei generi *Belonopelta* Mayr e *Leiopelta* gen. n. (Hymenoptera: Formicidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 48:295–310.
- _____. 1977. Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel. Hymenoptera: Fam. Formicidae Genus *Mayriella*. *Entomologica Basiliensis* 2:411–414.
- _____. 1978a. Contributo alla conoscenza del genere *Amblyopone* Erichson (Hymenoptera: Formicidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 51:39–51.
- _____. 1978b. Materiali per una revisione dei *Leptocephalus* neotropicali appartenenti al sotto-genere *Macromischus* Roger, n. comb. (Hymenoptera: Formicidae). *Entomologica Basiliensis* 3:395–618.
- Baroni Urbani, C., and M. L. De Andrade. 1993. *Perissomyrmex monticola* n. sp., from Bhutan: The first natural record for a presumed Neotropical genus with a discussion on its taxonomic status. *Tropical Zoology* 6:89–95.
- Baroni Urbani, C., B. Bolton, and P. S. Ward. 1992. The internal phylogeny of ants (Hymenoptera: Formicidae). *Systematic Entomology* 17:301–329.
- Barrer, P. M., and J. M. Cherrett. 1972. Some factors affecting the site and pattern of leaf-cutting activity in the ant *Atta cephalotes* L. *Journal of Entomology* 47:15–27.
- Basset, Y., N. D. Springate, H. P. Aberlenc, and G. Delvare. 1997. A review of methods for sampling arthropods in tree canopies. Pp. 27–52. In N. Stork, J. Adis, and R. K. Didham (eds.), *Canopy Arthropods*. Chapman and Hall, London.
- Beattie, A. J. 1985. The Evolutionary Ecology of Ant-Plant Mutualisms. Cambridge University Press, New York.
- Beattie, A., and I. Oliver. 1994. Taxonomic minimalism. *Trends in Evolution and Ecology* 9:488–490.
- Beattie, A. J., C. Turnbull, R. B. Knox, and E. G. Williams. 1984. Ant inhibition of pollen function: A possible reason why ant pollination is rare. *American Journal of Botany* 71:421–426.
- Beattie, A. J., C. Turnbull, T. Hough, S. Jobson, and R. B. Knox. 1985. The vulnerability of pollen and fungal spores to ant secretions: Evidence and some evolutionary implications. *American Journal of Botany* 72:606–614.
- Beattie, A. J., C. L. Turnbull, T. Hough, and R. B. Knox. 1986. Antibiotic production: A possible function for the metapleural glands of ants (Hymenoptera: Formicidae). *Annals of the Entomological Society of America* 79:448–450.
- Beccaloni, G. W., and K. J. Gaston. 1995. Predicting the richness of Neotropical forest butterflies: Ithomiinae (Lepidoptera: Nymphalidae) as indicators. *Biological Conservation* 71:77–86.
- Belshaw, R., and B. Bolton. 1993. The effect of forest disturbance on leaf litter ant fauna in Ghana. *Biodiversity and Conservation* 2:656–666.
- _____. 1994a. A survey of the leaf litter ant fauna in Ghana, West Africa (Hymenoptera: Formicidae). *Journal of Hymenoptera Research* 3:5–16.

- _____. 1994b. A new myrmicine ant genus from cocoa leaf litter in Ghana (Hymenoptera: Formicidae). *Journal of Natural History* 28:631–634.
- Belt, T. 1874. *The Naturalist in Nicaragua*. John Murray, London.
- Benson, W. W., and A. Y. Harada. 1988. Local diversity of tropical and temperate ant faunas. *Acta Amazonica* 18:275–289.
- Bequaert, J. 1922. Ants in their diverse relations to the plant world. *Bulletin of the American Museum of Natural History* 45:333–584, plates 26–29.
- Bernard, F. 1954. Fourmis moissonneuses nouvelles ou peu connues des montagnes d'Algérie et révision des *Messor* du groupe *structor* (Latr.). *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord* 45:354–365.
- _____. 1956. Révision des *Leptocephalus* (Hyménoptères Formicidae) d'Europe occidentale, basée sur la biométrie et les genitalia mâles. *Bulletin de la Société Zoologique de France* 81:151–165.
- _____. 1979. *Messor carthaginensis* n. sp., de Tunis, et révision des *Messor* du groupe *barbara* (Hym. Formicidae). *Bulletin de la Société Entomologique de France* 84:265–269.
- Bernstein, R. A. 1979. Schedules of foraging activity in species of ants. *Journal of Animal Ecology* 48:921–930.
- Bernstein, R. A., and M. Gobbel. 1979. Partitioning of space in communities of ants. *Journal of Animal Ecology* 48:931–942.
- Beschers, S., and J. Traniello. 1994. The adaptiveness of worker demography in the attine ant *Trachymyrmex septentrionalis*. *Ecology* 75:763–775.
- Bestelmeyer, B. T. 1997. Stress tolerance in some Chacoan dolichoderine ants: Implications for community organization and distribution. *Journal of Arid Environments* 35:297–310.
- Bestelmeyer, B. T., and J. A. Wiens. 1996. The effects of land use on the structure of ground-foraging ant communities in the Argentine chaco. *Ecological Applications* 6:1225–1240.
- Besuchet, C., D. H. Burckhardt, and I. Löbl. 1987. The "Winkler/Moczarski" elector as an efficient extractor for fungus and litter coleoptera. *The Coleopterists' Bulletin* 41:392–394.
- Billen, J. P. J. 1990. Phylogenetic aspects of exocrine gland development in the Formicidae. Pp. 317–318. In G. K. Veeresh, B. Mallik, and C. A. Viraktamath (eds.), *Social Insects and the Environment*. Proceedings of the 11th International Congress of IUSSI, 1990. Oxford and IBH, New Delhi.
- Bingham, C. T. 1903. The fauna of British India, including Ceylon and Burma. In *Hymenoptera*, Vol. II. Ants and Cuckoo-Wasps. Taylor and Francis, London.
- Black, R. W., II. 1987. The biology of leaf nesting ants in a tropical wet forest. *Biotropica* 19:319–325.
- Blackburn, T., P. Harvey, and M. Pagel. 1990. Species number, population density and body size relationships in natural communities. *Journal of Animal Ecology* 59:335–345.
- Bolton, B. 1972. Two new species of the ant genus *Epitritus* from Ghana, with a key to the world species (Hym., Formicidae). *Entomologists' Monthly Magazine* 107:205–208.
- _____. 1973a. A remarkable new arboreal ant genus (Hym. Formicidae) from West Africa. *Entomologists' Monthly Magazine* 108:234–237.
- _____. 1973b. The ant genus *Polyrhachis* F. Smith in the Ethiopian region (Hymenoptera: Formicidae). *Bulletin of the British Museum (Natural History)*. Entomology 28:283–369.
- _____. 1973c. New synonymy and a new name in the ant genus *Polyrhachis* F. Smith (Hym., Formicidae). *Entomologists' Monthly Magazine* 109:172–180.
- _____. 1974a. A revision of the Palaeotropical arboreal ant genus *Cataulacus* F. Smith (Hymenoptera: Formicidae). *Bulletin of the British Museum (Natural History)*. Entomology 30:1–105.
- _____. 1974b. A revision of the ponerine ant genus *Plectroctena* F. Smith (Hymenoptera: Formicidae). *Bulletin of the British Museum (Natural History)*. Entomology 30:309–338.
- _____. 1975a. A revision of the ant genus *Leptogenys* Roger (Hymenoptera: Formicidae) in the Ethiopian region with a review of the Malagasy species. *Bulletin of the British Museum (Natural History)*. Entomology 31:235–305.
- _____. 1975b. A revision of the African ponerine ant genus *Psalidomyrmex* André (Hymenoptera: Formicidae). *Bulletin of the British Museum (Natural History)*. Entomology 32:1–16.
- _____. 1975c. The *sexspinosa*-group of the ant genus *Polyrhachis* F. Smith (Hym. Formicidae). *Journal of Entomology Series B* 44:1–14.
- _____. 1976. The ant tribe Tetramoriini (Hymenoptera: Formicidae). Constituent genera, review of smaller genera and revision of *Triglyphothrix*

- Forel. Bulletin of the British Museum (Natural History). Entomology 34:281–379.
- . 1977. The ant tribe Tetramoriini (Hymenoptera: Formicidae). The genus *Tetramorium* Mayr in the Oriental and Indo-Australian regions, and in Australia. Bulletin of the British Museum (Natural History). Entomology 36:67–151.
- . 1979. The ant tribe Tetramoriini (Hymenoptera: Formicidae). The genus *Tetramorium* Mayr in the Malagasy region and in the New World. Bulletin of the British Museum (Natural History). Entomology 38:129–181.
- . 1980. The ant tribe Tetramoriini (Hymenoptera: Formicidae). The genus *Tetramorium* Mayr in the Ethiopian zoogeographical region. Bulletin of the British Museum (Natural History). Entomology 40:193–384.
- . 1981a. A revision of the ant genera *Meranoplus* F. Smith, *Dicroaspis* Emery and *Calyptomyrmex* Emery (Hymenoptera: Formicidae) in the Ethiopian zoogeographical region. Bulletin of the British Museum (Natural History). Entomology 42:43–81.
- . 1981b. A revision of six minor genera of Myrmicinae (Hymenoptera: Formicidae) in the Ethiopian zoogeographical region. Bulletin of the British Museum (Natural History). Entomology 43:245–307.
- . 1982. Afrotropical species of the myrmicine ant genera *Cardiocondyla*, *Leptothorax*, *Melissotarsus*, *Messor* and *Cataulacus* (Formicidae). Bulletin of the British Museum (Natural History). Entomology 45:307–370.
- . 1983. The Afrotropical dacetine ants (Formicidae). Bulletin of the British Museum (Natural History). Entomology 46:267–416.
- . 1984. Diagnosis and relationships of the myrmicine ant genus *Ishakidris* gen. n. (Hymenoptera: Formicidae). Systematic Entomology 9:373–382.
- . 1986. A taxonomic and biological review of the tetramoriine ant genus *Rhopromyrmex* (Hymenoptera: Formicidae). Systematic Entomology 11:1–17.
- . 1987. A review of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr (Hymenoptera: Formicidae). Bulletin of the British Museum (Natural History). Entomology 54:263–452.
- . 1988a. *Secostruma*, a new subterranean tetramoriine ant genus (Hymenoptera: Formicidae). Systematic Entomology 13:263–270.
- . 1988b. A review of *Paratopula* Wheeler, a forgotten genus of myrmicine ants (Hym., Formi-
- cidae). Entomologists' Monthly Magazine 124: 125–143.
- . 1990a. Abdominal characters and status of the cerapachyne ants (Hymenoptera, Formicidae). Journal of Natural History 24:53–68.
- . 1990b. [Untitled. A key to the living subfamilies of ants, based on the worker caste.] Pp. 33–34. In B. Hölldobler and E. O. Wilson, *The Ants*. Harvard University Press, Cambridge, Massachusetts.
- . 1990c. Army ants reassessed: The phylogeny and classification of the doryline section (Hymenoptera, Formicidae). Journal of Natural History 24:1339–1364.
- . 1990d. The higher classification of the ant subfamily Leptanillinae (Hymenoptera: Formicidae). Systematic Entomology 15:267–282.
- . 1991. New myrmicine genera from the Oriental Region (Hymenoptera: Formicidae). Systematic Entomology 16:1–13.
- . 1992. A review of the ant genus *Recurvidris* (Hym.: Formicidae), a new name for *Trigono-gaster* Forel. Psyche (Cambridge) 99:35–48.
- . 1994. Identification Guide to the Ant Genera of the World. Harvard University Press, Cambridge, Massachusetts.
- . 1995a. A taxonomic and zoogeographical census of the extant ant taxa (Hymenoptera: Formicidae). Journal of Natural History 29:1037–1056.
- . 1995b. A New General Catalogue of the Ants of the World. Harvard University Press, Cambridge, Massachusetts.
- . 1999. Ant genera of the tribe Dacetonini (Hymenoptera: Formicidae). Journal of Natural History 33:1639–1689.
- Bolton, B., and R. Belshaw. 1993. Taxonomy and biology of the supposedly lestobiotic ant genus *Paedalgus* (Hymenoptera: Formicidae). Systematic Entomology 18:181–189.
- Bolton, B., and A. C. Marsh. 1989. The Afrotropical thermophilic ant genus *Ocymyrmex* (Hymenoptera: Formicidae). Journal of Natural History 23:1267–1308.
- Bond, W., and P. Slingsby. 1984. Collapse of an ant-plant mutualism: The Argentine ant (*Iridomyrmex humilis*) and myrmecochorous proteaceae. Ecology 65:1031–1037.
- Bonham, C. D. 1989. Measurements for Terrestrial Vegetation. John Wiley and Sons, New York.
- Borgmeier, T. 1955. Die Wanderameisen der neotropischen Region. Studia Entomologica 3: 1–720.

- . 1959. Revision der Gattung *Atta* Fabricius (Hymenoptera, Formicidae). *Studia Entomologica* (n.s.) 2:321–390.
- . 1963. Revision of the North American phorid flies. Part I. The Phorinae, Aenigmatiinae, and Metopininae, except Megaselia (Diptera: Phoridae). *Studia Entomologica* (n.s.) 6:1–256.
- Borror, D., C. Triplehorn, and N. Johnson. 1989. An Introduction to the Study of Insects. W. B. Saunders, Philadelphia.
- Brandão, C. R. F. 1989. *Belonopelta minima*, a new species from Brazil. *Revista Brasileira de Entomologia* 33:135–138.
- . 1990. Systematic revision of the neotropical ant genus *Megalomyrmex* Forel (Hymenoptera: Formicidae: Myrmicinae) with the description of thirteen new species. *Arquivos de Zoologia (Museu de Zoologia da Universidade de São Paulo)* 31:411–481.
- . 1991. Adendos ao Catálogo Abreviado das Formigas da Região Neotropical (Hymenoptera: Formicidae). *Revista Brasileira de Entomologia* 35(2):319–412.
- Brandão, C. R. F., and J. E. Lattke. 1990. Description of a new Ecuadorian *Gnamptogenys* species (Hymenoptera: Formicidae), with a discussion on the status of the *alfaria* group. *Journal of the New York Entomological Society* 98:489–494.
- Brandão, C. R. F., and R. V. S. Paiva. 1994. The Galapagos ant fauna and the attributes of colonizing ant species. Pp. 1–10. In D. F. Williams (ed.), Exotic Ants: Biology, Impact, and Control of Introduced Species. Westview Press, Boulder, Colorado.
- Brandão, C. R. F., and P. E. Vanzolini. 1985. Notes on incubatory inquilinism between squamata (Reptilia) and the neotropical fungus-growing ant genus *Acromyrmex* (Hymenoptera: Formicidae). *Papeis Avulsos de Zoologia (São Paulo)* 36:31–36.
- Brandão, C. R. F., J. L. M. Diniz, and E. M. Tomotake. 1991. *Thaumatomyrmex* strip millipedes for prey: A novel predatory behaviour in ants, and the first case of sympatry in the genus (Hymenoptera: Formicidae). *Insectes Sociaux* 38:335–344.
- Brandão, C. R. F., J. L. M. Diniz, D. Agosti, and J. H. Delabie. 1999. Revision of the Neotropical ant subfamily Leptanilloidinae. *Systematic Entomology* 24:17–36.
- Brian, M. V. 1964. Ant distribution in a southern English heath. *Journal of Animal Ecology* 33:451–461.
- Brian, M. V., and A. D. Brian. 1951. Insolation and ant populations in the west of Scotland. *Transactions of the Royal Entomological Society of London* 102:303–330.
- Brian, M. V., M. D. Mountford, A. Abbott, and S. Vincent. 1976. The changes in ant species distribution during ten years' post-fire regeneration of a heath. *Journal of Animal Ecology* 45:115–133.
- Brothers, D. J. 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutilidae. *University of Kansas Science Bulletin* 50:483–648.
- Brothers, D. J., and J. M. Carpenter. 1993. Phylogeny of Aculeata: Chrysidoidea and Vespoidea (Hymenoptera). *Journal of Hymenoptera Research* 2:227–304.
- Brown, B. V. 1993. Taxonomy and preliminary phylogeny of the parasitic genus *Apocephalus*, subgenus *Mesophora* (Diptera: Phoridae). *Systematic Entomology* 18:191–230.
- Brown, B. V., and D. H. Feener Jr. 1991a. Behavior and host location cues of *Apocephalus paraponerae* (Diptera: Phoridae), a parasitoid of the giant tropical ant, *Paraponera clavata* (Hymenoptera: Formicidae). *Biotropica* 23:182–187.
- . 1991b. Life history parameters and description of the larva of *Apocephalus paraponerae* (Diptera: Phoridae), a parasitoid of the giant tropical ant *Paraponera clavata* (Hymenoptera: Formicidae). *Journal of Natural History* 25:221–231.
- Brown, J. H. 1995. Macroecology. University of Chicago Press, Chicago.
- Brown, J. H., T. Valone, and C. Curtin. 1997. Reorganization of an arid ecosystem in response to recent climate change. *Proceedings of the National Academy of Sciences* 94:9729–9733.
- Brown, W. L., Jr. 1945. An unusual behavior pattern observed in a Szechuanese ant. *Journal of the West China Border Research Society, Series B* 15:185–186.
- . 1948. A preliminary generic revision of the higher Dacetini (Hymenoptera: Formicidae). *Transactions of the American Entomological Society* 74:101–129.
- . 1949a. Synonymic and other notes on Formicidae (Hymenoptera). *Psyche (Cambridge)* 56:41–49.
- . 1949b. Revision of the ant tribe Dacetini. III. *Epitritus* Emery and *Quadristruma* new genus (Hymenoptera: Formicidae). *Transactions of the American Entomological Society* 75:43–51.

- _____. 1949c. Revision of the ant tribe Dacetini. I. Fauna of Japan, China and Taiwan. Mushi 20: 1–25.
- _____. 1949d. Revision of the ant tribe Dacetini. IV. Some genera properly excluded from the Dacetini, with the establishment of the Basicerotini new tribe. Transactions of the American Entomological Society 75:83–96.
- _____. 1950. Revision of the ant tribe Dacetini. II. *Glamyromyrmex* Wheeler and closely related small genera. Transactions of the American Entomological Society 76:27–36.
- _____. 1952. On the identity of *Adlerzia* Forel (Hymenoptera: Formicidae). Pan-Pacific Entomologist 28:173–177.
- _____. 1953a. Revisionary studies in the ant tribe Dacetini. American Midland Naturalist 50:1–137.
- _____. 1953b. Characters and synonymies among the genera of ants. Part II. Breviora 18:1–8.
- _____. 1953c. Revisionary notes on the ant genus *Myrmecia* of Australia. Bulletin of the Museum of Comparative Zoology, Harvard University 111:1–35.
- _____. 1954a. (1953) The Indo-Australian species of the ant genus *Strumigenys* Fr. Smith: S. wallacei Emery and relatives. Psyche (Cambridge) 60:85–89.
- Brown, W. L., Jr. 1954b. Remarks on the internal phylogeny and subfamily classification of the family Formicidae. Insectes Sociaux 1:22–31.
- _____. 1955. A revision of the Australian ant genus *Notoncus* Emery, with notes on the other genera of Melophorini. Bulletin of the Museum of Comparative Zoology, Harvard College 113:471–494.
- _____. 1958. Contributions toward a reclassification of the Formicidae. II. Tribe Ectatommini (Hymenoptera). Bulletin of the Museum of Comparative Zoology, Harvard University 118:173–362.
- _____. 1959. A revision of the dacetine ant genus *Neostruma*. Breviora 107:1–13.
- _____. 1960. Contributions toward a reclassification of the Formicidae. III. Tribe Amblyoponini (Hymenoptera). Bulletin of the Museum of Comparative Zoology, Harvard University 122:143–230.
- _____. 1962. The neotropical species of the ant genus *Strumigenys* Fr. Smith: Synopsis and keys to the species. Psyche (Cambridge) 69:238–267.
- _____. 1964. The ant genus *Smithistruma*: A first supplement to the World revision (Hymenoptera: Formicidae). Transactions of the American Entomological Society 89:183–200.
- _____. 1965. Contributions to a reclassification of the Formicidae. IV. Tribe Typhlomyrmecini (Hymenoptera). Psyche (Cambridge) 72:65–78.
- _____. 1967. Studies on North American ants. II. *Myrmecina*. Entomological News 78:233–240.
- _____. 1972. *Aasketogenys acubbecca*, a new genus and species of dacetine ants from Malaya (Hymenoptera: Formicidae). Psyche (Cambridge) 79:23–26.
- _____. 1973. A comparison of the Hylean and Congo-West African rain forest ant faunas. Pp. 161–185. In B. J. Meggers, E. S. Ayensu, and W. D. Duckworth (eds.), Tropical Forest Ecosystems in Africa and South America: A Comparative Review. Smithsonian Institution Press, Washington, D.C.
- _____. 1974a. A supplement to the revision of the ant genus *Basiceros* (Hymenoptera: Formicidae). Journal of the New York Entomological Society 82:131–140.
- _____. 1974b. *Concoctio* genus nov. Pilot Register of Zoology. Card No. 29.
- _____. 1974c. *Concoctio concentra* species nov. Pilot Register of Zoology. Card No. 30.
- _____. 1974d. *Dolioponera* genus nov. Pilot Register of Zoology. Card No. 31.
- _____. 1974e. *Dolioponera fustigera* species nov. Pilot Register of Zoology. Card No. 32.
- _____. 1975. Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythyreini, Cerapachyini, Cylindromyrmecini, Acanthostichini, and Aenictogitini. Search Agriculture (Ithaca, N.Y.) 5(1):1–115.
- _____. 1976a. *Cladarogenys* genus nov. Pilot Register of Zoology. Card No. 33.
- _____. 1976b. *Cladarogenys lasia* species nov. Pilot Register of Zoology. Card No. 34.
- _____. 1976c. Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section A. Introduction, subtribal characters. Genus *Odontomachus*. Studia Entomologica 19:67–171.
- _____. 1977a. An aberrant new genus of myrmicine ant from Madagascar. Psyche (Cambridge) 84:218–224.
- _____. 1977b. A supplement to the world revision of *Odontomachus* (Hymenoptera: Formicidae). Psyche (Cambridge) 84:281–285.
- _____. 1978. Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe

- Ponerini, subtribe Odontomachiti. Section B. Genus *Anochetus* and bibliography. *Studia Entomologica* 20:549–638.
- . 1979. A remarkable new species of *Proceratium*, with dietary and other notes on the genus (Hymenoptera: Formicidae). *Psyche* (Cambridge) 86:337–346.
- . 1980a. *Protalaridris* genus nov. Pilot Register of Zoology. Card No. 36.
- . 1980b. *Protalaridris armata* species nov. Pilot Register of Zoology. Card No. 37.
- . 1985. *Indomyrma dasypyx*, new genus and species, a myrmicine ant from peninsular India (Hymenoptera: Formicidae). *Israel Journal of Entomology* 19:37–49.
- Brown, W. L., Jr., and R. G. Boisvert. 1979. The dacetine ant genus *Pentastrum*. *Psyche* (Cambridge) 85:201–207.
- Brown, W. L., Jr., and W. W. Kempf. 1960. A world revision of the ant tribe Basicerotini. *Studia Entomologica* (n.s.) 3:161–250.
- . 1967. *Tatuidris*, a remarkable new genus of Formicidae (Hymenoptera). *Psyche* (Cambridge) 74:183–190.
- . 1969. A revision of the neotropical dacetine ant genus *Acanthognathus* (Hymenoptera: Formicidae). *Psyche* (Cambridge) 76:87–109.
- Brown, W. L., Jr., W. H. Gotwald Jr., and J. Lévieux. 1970. A new genus of ponerine ants from West Africa (Hymenoptera: Formicidae) with ecological notes. *Psyche* (Cambridge) 77:259–275.
- Brues, C. T. 1925. *Scyphodon*, an anomalous genus of Hymenoptera of doubtful affinities. *Treubia* 6:93–96.
- Buckley, R. C. 1982a. Ant-plant interactions: A world review. Pp. 111–141. In R. C. Buckley (ed.), *Ant-Plant Interactions in Australia*. Dr. W. Junk, The Hague.
- . 1982b. A world bibliography of ant-plant interactions. Pp. 143–162. In R. C. Buckley (ed.), *Ant-Plant Interactions in Australia*. Dr. W. Junk, The Hague.
- Bukowski, T. C. 1991. Solifugae in *Atta* foraging columns. P. 70. In B. A. Loiselle and C. K. Augspurger (eds.), OTS 91-1: Tropical Biology: An Ecological Approach. Organization for Tropical Studies, Duke University, Durham, North Carolina.
- Bunge, J., and M. Fitzpatrick. 1993. Estimating the number of species: A review. *Journal of the American Statistical Association* 88:364–373.
- Bünzli, G. H. 1935. Untersuchungen über coccophile Ameisen aus den Kaffeefeldern von Surinam. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 16:453–593.
- Burbridge, A. H., K. Leicester, S. McDavitt, and J. D. Majer. 1992. Ants as indicators of disturbance at Yanchep National Park, Western Australia. *Journal of the Royal Society of Western Australia* 75:89–95.
- Buren, W. F. 1968a. Some fundamental taxonomic problems in *Formica* (Hymenoptera: Formicidae). *Journal of the Georgia Entomological Society* 3:25–40.
- . 1968b. A review of the species of *Crematogaster*, sensu stricto, in North America (Hymenoptera, Formicidae). Part II. Descriptions of new species. *Journal of the Georgia Entomological Society* 3:91–121.
- Burnham, K. P., and W. S. Overton. 1978. Estimation of the size of a closed population when capture probabilities vary among animals. *Biometrika* 65:623–633.
- . 1979. Robust estimation of population size when capture probabilities vary among animals. *Ecology* 60:927–936.
- Buschinger, A. 1981. Biological and systematic relationships of social parasitic Leptocephalacini from Europe and North America. Pp. 211–222. In P. E. Howse and J.-L. Clement (eds.), *Biosystematics of Social Insects*. Systematics Association Special Volume No. 19. Academic Press, London.
- . 1989. Evolution, speciation, and inbreeding in the parasitic ant genus *Epimyrma* (Hymenoptera, Formicidae). *Journal of Evolutionary Biology* 2:265–283.
- Buschinger, A., and U. Winter. 1983. *Myrmicinosporidium durum* Hölldobler 1933, Parasit bei Ameisen (Hym., Formicidae) in Frankreich, der Schweiz und Jugoslawien wieder aufgefunden. *Zoologischer Anzeiger* 210:393–398.
- Buschinger, A., J. Heinze, K. Jessen, P. Douwes, and U. Winter. 1987. First European record of a queen ant carrying a mealybug during her mating flight. *Naturwissenschaften* 74:139–140.
- Buschinger, A., W. Ehrhardt, K. Fischer, and J. Ofer. 1988. The slave-making ant genus *Chalepoxenus* (Hymenoptera, Formicidae). I. Review of literature, range, slave species. *Zoologische Jahrbücher. Abteilung für Systematik, Ökologie und Geographie der Tiere* 115:383–401.

- Buschinger, A., R. G. Kleespies, and R. D. Schumann. 1995. A gregarine parasite of *Leptothorax* ants from North America. *Insectes Sociaux* 42:219–222.
- Byrne, M. M. 1994. Ecology of twig-dwelling ants in a wet lowland tropical forest. *Biotropica* 26:61–72.
- Caetano, F. H. 1989. Endosymbiosis of ants with intestinal and salivary gland bacteria. In W. Schwemmle and G. Gassner (eds.), *Insect Endosymbionts: Morphology, Physiology, Genetics, Evolution*. CRC Press, Boca Raton, Florida.
- Caetano, F. H., and C. Cruz-Landim. 1985. Presence of microorganisms in the alimentary canal of ants of the tribe Cephalotini (Myrmicinae): Location and relationship with intestinal structures. *Naturalia* 10:37–47.
- Cagniant, H. 1997. The ant genus *Tetramorium* (Hymenoptera: Formicidae) in Morocco. *Annales de la Société Entomologique de France* 33(1):89–100.
- Cagniant, H., and X. Espadaler. 1997a. *Leptothorax, Epimyrma* and *Chalepoxenus* of Morocco (Hymenoptera: Formicidae). Key and catalogue of species. *Annales de la Société Entomologique de France* 33(3):259–284.
- Cagniant, H., and X. Espadaler. 1997b. The ant genus *Messor* in Morocco (Hymenoptera: Formicidae). *Annales de la Société Entomologique de France* 33(4):419–434.
- Camilo, G. R., and S. A. Phillips Jr. 1990. Evolution of ant communities in response to invasion by the fire ant *Solenopsis invicta*. Pp. 190–198. In R. K. Vander Meer, K. Jaffe, and A. Cedeno (eds.), *Applied Myrmecology: A World Perspective*. Westview Press, Boulder, Colorado.
- Cammell, M. E., M. J. Way, and M. R. Paiva. 1996. Diversity and structure of ant communities associated with oak, pine, eucalyptus, and arable habitats in Portugal. *Insectes Sociaux* 43:37–46.
- Carpenter, S., T. Frost, J. Kitchell, and T. Kratz. 1993. Species dynamics and global environmental change: A perspective from ecosystem experiments. Pp. 267–279. In P. Kareiva, J. Kingsolver, and R. Huey (eds.), *Biotic Interactions and Global Change*. Sinauer Associates, Sunderland, Massachusetts.
- Chao, A. 1984. Non-parametric estimation of the number of classes in a population. *Scandinavian Journal of Statistics* 11:265–270.
- . 1987. Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* 43:783–791.
- Chao, A., and S. M. Lee. 1992. Estimating the number of classes via sample coverage. *Journal of the American Statistical Association* 87:210–217.
- Chao, A., M. C. Ma, and M. C. K. Yang. 1993. Stopping rules and estimation for recapture debugging with unequal failure rates. *Biometrika* 80:193–201.
- Chapela, I. H., S. A. Rehner, T. R. Schultz, and U. G. Mueller. 1994. Evolutionary history of the symbiosis between the fungus-growing ants and their fungi. *Science* 266:1691–1694.
- Chapman, T. A. 1920. Contributions to the life history of *Lycaena euphemus* Hb. *Transactions of the Royal Entomological Society of London* 1919:450–465.
- Chazdon, R. L., R. K. Colwell, J. S. Denslow, and M. R. Guariguata. 1998. Statistical methods for estimating species richness of woody regeneration in primary and secondary rain forests of northeastern Costa Rica. Pp. 285–309. In F. Dallmeier and J. A. Comiskey (eds.), *Forest Biodiversity Research, Monitoring and Modeling: Conceptual Background and Old World Case Studies*. Parthenon, Paris.
- Chen, Y.-C., W.-H. Hwang, A. Chao, and C.-Y. Kuo. 1995. Estimating the number of common species. Analysis of the number of common bird species in Ke-Yar Stream and Chung-Kang Stream. *Journal of the Chinese Statistical Association* 33:373–393. [In Chinese with English abstract.]
- Cherrett, J. M. 1986. History of the leaf-cutting ant problem. Pp. 10–17. In C. S. Lofgren and R. K. Vander Meer (eds.), *Fire Ants and Leaf Cutting Ants: Biology and Management*. Westview Press, Boulder, Colorado.
- Chew, R. M. 1995. Aspects of the ecology of three species of ants (*Myrmecocystus* spp., *Aphaenogaster* sp.) in desertified grassland in southeastern Arizona, 1958–1993. *American Midland Naturalist* 134:75–83.
- Chew, R. M., and J. De Vita. 1980. Foraging characteristics of a desert ant assemblage: Functional morphology and species separation in Cochise County, Arizona. *Journal of Arid Environments* 3:75–83.
- Christian, K., and S. R. Morton. 1992. Extreme thermophilia in a central Australian ant, *Melophorus bagoti*. *Physiological Zoology* 65:885–905.
- Clark, D. B., C. Guayasamín, O. Pazmiño, C. Donoso, and Y. Páez de Villacís. 1982. The tramp ant *Wasmannia auropunctata*: Autecology and effects

- on ant diversity and distribution on Santa Cruz Island, Galapagos. *Biotropica* 14:196–207.
- Clark, J. 1930. The Australian ants of the genus *Dolichoderus* (Formicidae). Subgenus *Hypoclinea* Mayr. *Australian Zoologist* 6:252–268.
- . 1936. A revision of Australian species of *Rhytidoponera* Mayr (Formicidae). *Memoirs of the National Museum of Victoria* 9:14–89.
- . 1951. The Formicidae of Australia, Vol. 1: Subfamily Myrmeciinae. CSIRO, Melbourne.
- Clausen, C. P. 1940a. *Entomophagous Insects*. McGraw-Hill, New York.
- . 1940b. The immature stages of the Eucharidae. *Proceedings of the Entomological Society of Washington* 42:161–170.
- . 1940c. The oviposition habits of the Eucharidae (Hymenoptera). *Journal of the Washington Academy of Sciences* 30:504–516.
- . 1941. The habits of the Eucharidae. *Psyche (Cambridge)* 48:57–69.
- Clements, R. O. 1982. Sampling and extraction techniques for collecting invertebrates from grassland. *Entomologists' Monthly Magazine* 118:133–142.
- Clench, H. K. 1979. How to make regional lists of butterflies: Some thoughts. *Journal of the Lepidopterist Society* 33:216–231.
- Cobabe, E. A., and W. D. Allmon. 1994. Effects of sampling on paleoecologic and taphonomic analyses in high-diversity fossil accumulations: An example from the Eocene Gosport Sand, Alabama. *Lethaia* 27:167–178.
- Coddington, J. A., L. H. Young, and F. A. Coyle. 1996. Estimating spider species richness in a southern Appalachian cove hardwood forest. *Journal of Arachnology* 24:111–128.
- Coenen-Stass D., Schaarschmidt, and I. Lamprecht. 1980. Temperature distribution and calorimetric determination of heat production in the nest of the wood ant, *Formica polyctena* (Hymenoptera: Formicidae). *Ecology* 61:238–244.
- Cole, A. C., Jr. 1940. A guide to the ants of the Great Smoky Mountains National Park, Tennessee. *American Midland Naturalist* 24:1–88.
- . 1949. Notes on *Gesomyrmex* (Hymenoptera: Formicidae). *Entomological News* 60:181.
- . 1968. *Pogonomyrmex* Harvester Ants: A Study of the Genus in North America. University of Tennessee Press, Knoxville.
- Colwell, R. K. 1997. EstimateS: Statistical estimation of species richness and shared species from samples. Version 5. User's guide and application published at: <http://viceroy.eeb.uconn.edu/> estimates.
- Colwell, R. K., and J. A. Coddington. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society of London, Series B* 345:101–118.
- Colwell, R. K., and G. C. Hurtt. 1994. Nonbiological gradients in species richness and a spurious Rapaport effect. *American Naturalist* 144:570–595.
- Connell, J. 1978. Diversity in tropical rain forests and coral reefs. *Science* 199:1302–1310.
- Connell, J., and W. Sousa. 1983. On the evidence needed to judge ecological stability or persistence. *American Naturalist* 121:789–824.
- Convention on Biological Diversity. 1992. Convention on Biological Diversity. United Nations Conference on Environment and Development, Rio de Janeiro.
- Cover, S. P., J. E. Tobin, and E. O. Wilson. 1990. The ant community of a tropical lowland rain forest site in Peruvian Amazonia. Pp. 699–700. In G. K. Veeresh, B. Mallik, and C. A. Viraktamath (eds.), *Social Insects and the Environment*. Proceedings of the 11th International Congress of IUSSI, 1990. Oxford and IBH Publishing, New Delhi.
- Cowling, R. M., and J. J. Midgely. 1996. The influence of regional phenomena on an emerging global ecology. *Global Ecology and Biogeography Letters* 5:63–65.
- Cranston, P. S., and J. W. H. Trueman. 1997. "Indicator" taxa in invertebrate biodiversity assessment. *Memoirs of the Museum of Victoria* 56(2):267–274.
- Creighton, W. S. 1950. The ants of North America. *Bulletin of the Museum of Comparative Zoology, Harvard College* 104:1–585.
- . 1957. A study of the genus *Xenomyrmex* (Hymenoptera, Formicidae). *American Museum Novitates* 1843:1–14.
- Crist, T. O., and J. A. Wiens. 1994. Scale effects of vegetation structure on forager movements and seed harvesting by ants. *Oikos* 69:37–46.
- . 1996. The distribution of ant colonies in a semiarid landscape: Implications for community and ecosystem processes. *Oikos* 76:301–311.
- Crosland, M. W. J. 1988. Effect of a gregarine parasite on the color of *Myrmecia pilosula* (Hymenoptera: Formicidae). *Annals of the Entomological Society of America* 81:481–484.

- Crowell, K. L. 1968. Rates of competitive exclusion by the Argentine ant in Bermuda. *Ecology* 49:551–555.
- Crozier, R. H. 1990. From population genetics to phylogeny: Uses and limits of mitochondrial DNA. *Australian Systematic Botany* 3:111–124.
- Culver, D. S. 1974. Species packing in Caribbean and North temperate ant communities. *Ecology* 55:974–988.
- Cushman, J. H., J. H. Lawton, and B. F. J. Manly. 1993. Latitudinal patterns in European ant assemblages: Variation in species richness and body size. *Oecologia* 95:30–37.
- Darlington, P. J., Jr. 1971. The carabid beetles of New Guinea. Part IV. General considerations; analysis and history of fauna; taxonomic supplement. *Bulletin of the Museum of Comparative Zoology, Harvard College* 142:129–337.
- Davidson, D. W. 1977a. Species diversity and community organization in desert seed-eating ants. *Ecology* 58:711–724.
- . 1977b. Foraging ecology and community organization in desert seed-eating ants. *Ecology* 58:725–737.
- . 1978. Size variability in the worker caste of a social insect (*Veromessor pergandei* Mayr) as a function of the competitive environment. *American Naturalist* 112:523–532.
- . 1980. Some consequences of diffuse competition in a desert ant community. *American Naturalist* 116:92–105.
- . 1988. Ecological studies of Neotropical ant gardens. *Ecology* 69:1138–1152.
- . 1997. The role of resource imbalances in the evolutionary ecology of tropical arboreal ants. *Biological Journal of the Linnean Society* 61:153–181.
- . 1998. Resource discovery versus resource domination in ants: A functional mechanism for breaking the tradeoff. *Ecological Entomology* 23:484–490.
- Davidson, D. W., and L. Patrell-Kim. 1996. Tropical arboreal ants: Why so abundant? Pp. 127–140. In A. C. Gibson (ed.), *Neotropical Biodiversity and Conservation*. University of California, Los Angeles Botanical Garden Publication Number 1. University of California, Los Angeles.
- Davidson, D. W., J. H. Brown, and R. S. Inouye. 1980. Competition and the structure of granivore communities. *BioSciences* 30(4):233–238.
- Davidson, D. W., J. T. Longino, and R. R. Snelling. 1988. Pruning of host plant neighbors by ants: An experimental approach. *Ecology* 69:801–808.
- Davidson, D. W., R. R. Snelling, and J. T. Longino. 1989. Competition among ants for myrmecophytes and the significance of plant trichomes. *Biotropica* 21:64–73.
- De Kock, A. E., and J. H. Giliomee. 1989. A survey of the Argentine ant, *Iridomyrmex humilis* (Mayr) (Hymenoptera: Formicidae) in South African fynbos. *Journal of the Entomological Society of Southern Africa* 52:157–164.
- De Vries, P. J., D. Murray, and R. Lande. 1997. Species diversity in vertical, horizontal, and temporal dimensions of a fruit-feeding butterfly community in an Ecuadorian rainforest. *Biological Journal of the Linnean Society* 62:343–364.
- Delabie, J. H. C. 1995. Inquilinismo simultâneo de duas espécies de *Centromyrmex* (Hymenoptera; Formicidae; Ponerinae) em cupinzeiros de *Syntermes* sp. (Isoptera; Termitidae; Nasutitermitinae). *Revista Brasileira de Entomologia* 39:605–609.
- Delabie, J. H. C., and H. G. Fowler. 1995. Soil and litter cryptic ant assemblages of Bahian cocoa plantations. *Pedobiologia* 39:423–433.
- Delabie, J. H. C., I. C. do Nascimento, and C. dos S. F. Mariano. 2000. Importance de l'agriculture cacaoyère pour le mainien de la biodiversité: Étude comparée de la myrmécofaune de différents milieux du sud-est de Bahia, Brésil (Hymenoptera; Formicidae). In *Proceedings for the 12th International Cocoa Research Conference*, Lagos, Nigeria.
- Delabie, J. H. C., A. B. Casimiro, I. C. do Nascimento, A. L. B. do Souza, M. Furst, A. M. V. da Encarnação, M. R. B. Smith, and I. M. Cazorla. 1994. Stratification de la communauté de fourmis (Hymenoptera: Formicidae) dans une cacaoyère brésilienne et conséquences pour le contrôle naturel des ravageurs du cacaoyer. Pp. 823–831. In *Proceedings of the 11th International Cocoa Research Conference*, Lagos, Nigeria.
- Delabie, J. H. C., I. C. do Nascimento, P. Pacheco, and A. B. Casimiro. 1995. Community structure of house-infesting ants (Hymenoptera: Formicidae) in Southern Bahia, Brazil. *Florida Entomologist* 78:264–270.
- Deslippe, R. J., and R. Savolainen. 1994. Role of food supply in structuring a population of *Formica* ants. *Journal of Animal Ecology* 63:756–764.

- Deyrup, M., J. Trager, and N. Carlin. 1985. The genus *Odontomachus* in the southeastern United States (Hymenoptera: Formicidae). Entomological News 96:188–195.
- Di Castri, F., J. Robertson Vernhes, and T. Younes. 1992. Inventorying and monitoring biodiversity. Biology International. 27:1–27.
- Diniz, J. L. M. 1990. Revisão sistemática da tribo Stegomymicini, com a descrição de uma nova espécie (Hymenoptera, Formicidae). Revista Brasileira de Biologia 34:277–295.
- Diniz, J. L. M., and C. R. F. Brandão. 1993. Biology and myriapod egg predation by the Neotropical myrmicine ant *Stegomyrmex vizottoi* (Hymenoptera: Formicidae). Insectes Sociaux 40:301–311.
- Dixon, A. F. G. 1985. Aphid Ecology. Chapman and Hall, New York.
- Dlussky, G. M. 1964. The ants of the subgenus *Coptoformica* of the genus *Formica* (Hymenoptera, Formicidae) of the USSR. Zoologicheskii Zhurnal 43:1026–1040. [In Russian.]
- . 1965. Ants of the genus *Formica* L. of Mongolia and northeast Tibet (Hymenoptera, Formicidae). Annales Zoologici (Warsaw) 23:15–43.
- . 1967. Ants of the Genus *Formica* (Hymenoptera, Formicidae, g. *Formica*). Nauka, Moscow. [In Russian.]
- . 1969. Ants of the genus *Proformica* Ruzs. of the USSR and contiguous countries (Hymenoptera, Formicidae). Zoologicheskii Zhurnal 48: 218–232. [In Russian.]
- Dlussky, G. M., and B. Pisarski. 1971. Rewizja polskich gatunków mrówek (Hymenoptera: Formicidae) z rodzaju *Formica* L. Fragmenta Faunistica (Warsaw) 16:145–224.
- Dlussky, G. M., and A. G. Radchenko. 1994. Ants of the genus *Diplorhoptrum* (Hymenoptera, Formicidae) from the central Palearctic. Zoologicheskii Zhurnal 73(2):102–111. [In Russian.]
- Dlussky, G. M., and O. S. Soyunov. 1988. Ants of the genus *Temnothorax* Mayr (Hymenoptera: Formicidae) of the USSR. Izvestiya Akademii Nauk Turkmenskoi SSR, Seriya Biologicheskikh Nauk 1988(4):29–37. [In Russian.]
- Donisthorpe, H. 1946. *Ireneopone gibber* (Hym., Formicidae), a new genus and species of myrmicine ant from Mauritius. Entomologists' Monthly Magazine 82:242–243.
- Dorow, W. H. O., and R. J. Kohout. 1995. A review of the subgenus *Hemioptica* Roger of the genus *Polyrhachis* Fr. Smith with description of a new species (Hymenoptera: Formicidae: Formicinae). Zoologische Mededelingen (Leiden) 69(1–14): 93–104.
- DuBois, M. B. 1981. Two new species of inquiline *Monomorium* from North America (Hymenoptera: Formicidae). University of Kansas Science Bulletin 52:31–37.
- . 1986. A revision of the native New World species of the ant genus *Monomorium* (*minimum* group) (Hymenoptera: Formicidae). University of Kansas Science Bulletin 53:65–119.
- . 1998. A revision of the ant genus *Stenamma* in the Palaearctic and Oriental regions (Hymenoptera: Formicidae: Myrmicinae). Sociobiology 32(2):193–403.
- Dumpert, K. 1981. The Social Biology of Ants. Translated by C. Johnson. Pitman, Boston.
- . 1985. *Camponotus (Karavaievia) texens* sp. n. and *C. (K.) gombaki* sp. n. from Malaysia in comparison with other *Karavaievia* species (Formicidae: Formicinae). Psyche (Cambridge) 92:557–573.
- Dumpert, K., U. Maschwitz, A. Weissflog, K. Rosciszewski, and I. H. Azarae. 1995. Six new weaver ant species from Malaysia: *Camponotus (Karavaievia) striaticeps*, *C. (K.) melanurus*, *C. (K.) nigripes*, *C. (K.) belumensis*, *C. (K.) gentingensis*, and *C. (K.) micragyne*. Malaysian Journal of Science 16A:87–105.
- Eisner, T. 1957. A comparative morphological study of the proventriculus of ants (Hymenoptera: Formicidae). Bulletin of the Museum of Comparative Zoology, Harvard College 116:439–490.
- Elmes, G. W. 1991. Ant colonies and environmental disturbance. Pp. 15–32. In P. S. Meadows and A. Meadows (eds.), Environmental Impact of Burrowing Animals and Animal Burrows. Clarendon Press, Oxford.
- Emery, C. 1897. Revisione del genere *Diacamma* Mayr. Rendiconti delle Sessioni dell' Accademia delle Scienze dell' Istituto di Bologna (n.s.) 1:147–167.
- . 1901. Notes sur les sous-familles des Dorylines et Ponérines (Famille des Formicides). Annales de la Société Entomologique de Belgique 45:32–54.
- . 1910. Hymenoptera. Fam. Formicidae. Subfam. Dorylinae. Genera Insectorum 102:1–34.
- . 1911. Hymenoptera. Fam. Formicidae. Subfam. Ponerinae. Genera Insectorum 118:1–125.

- _____. 1913 (1912). Hymenoptera. Fam. Formicidae. Subfam. Dolichoderinae. Genera Insectorum 137:1–50.
- _____. 1920. La distribuzione geografica attuale delle formiche. Tentativo di spiegarne la genesi col soccorso di ipotesi filogenetiche e paleogeografiche. Atti della Reale Accademia dei Lincei. Memorie. Classe di Scienze, Fisiche, Matematiche e Naturali (5)13:357–450.
- _____. 1921. Hymenoptera. Fam. Formicidae. Subfam. Myrmicinae. [part] Genera Insectorum 174A:1–94 + 7 plates.
- _____. 1922. Hymenoptera. Fam. Formicidae. Subfam. Myrmicinae. [part] Genera Insectorum 174B:95–206.
- _____. 1925a. Hymenoptera. Fam. Formicidae. Subfam. Formicinae. Genera Insectorum 183:1–302.
- _____. 1925b. Revision des espèces paléarctiques du genre *Tapinoma*. Revue Suisse Zoologique 32:45–64.
- Entomological Society of Canada. 1978. Collections of Canadian Insects and Certain Related Groups. Entomological Society of Canada, Ottawa.
- Erickson, J. M. 1971. The displacement of native ant species by the introduced Argentine ant *Iridomyrmex humilis* Mayr. *Psyche* (Cambridge) 78:257–266.
- Erwin, T. L. 1983. Beetles and other arthropods of tropical forest canopies at Manaus, Brazil, sampled by insecticidal fogging. Pp. 59–79. In S. L. Sutton, T. C. Whitmore, and A. C. Chadwick (eds.), *Ecology and Management of Tropical Rainforest*. Blackwell, Oxford.
- _____. 1986. The tropical forest canopy: The heart of biotic diversity. Pp. 123–129. In E. O. Wilson (ed.), *Biodiversity*. National Academy Press, Washington, D.C.
- _____. 1989. Sorting tropical forest canopy samples (an experimental project for networking information). *Insect Collection News* 2(1):8.
- Espadaler, X. 1982. *Myrmicinosporidium* sp., parasite interne des fourmis: Etude au meb de la structure externe. Pp. 239–241. In A. deHaro and X. Espadaler (eds.), *La Communication chez les Sociétés d'Insectes. Colloque Internationale de l'Union Internationale pour l'Etude des Insectes Sociaux*, Section française, Barcelona, 1982. Universidad Autónoma de Barcelona, Bellaterra.
- Ettershank, G. 1966. A generic revision of the world Myrmicinae related to *Solenopsis* and *Pheidoloides* (Hymenoptera: Formicidae). *Australian Journal of Zoology* 14:73–171.
- Evans, H. E. 1962. A review of nesting behavior of digger wasps of the genus *Aphilanthops*, with special attention to the mechanics of prey carriage. *Behavior* 19:239–260.
- _____. 1977. Prey specificity in *Clypeadon* (Hymenoptera: Sphecidae). *Pan-Pacific Entomologist* 53:144.
- Farquharson, C. O. 1914. The growth of fungi on the shelters built over Coccidae by *Cremastogaster*-ants. *Transactions of the Entomological Society of London* 1914:42–50.
- _____. 1918. *Harpagomyia* and other Diptera fed by *Cremastogaster* ants in S. Nigeria. *Proceedings of the Entomological Society of London* 1918:29–39.
- Feener, D. H., Jr. 1981. Competition between ant species: Outcome controlled by parasitic flies. *Science* 214:815–817.
- Feener, D. H., Jr., and K. A. G. Moss. 1990. Defense against parasites by hitchhikers in leaf-cutting ants: A quantitative assessment. *Behavioral Ecology and Sociobiology* 26:17–29.
- Feener, D. H., Jr., L. F. Jacobs, and J. O. Schmidt. 1996. Specialized parasitoid attracted to a pheromone of ants. *Animal Behavior* 51:61–66.
- Fellers, G. M., and J. H. Fellers. 1982. Scavenging rates of invertebrates in an eastern deciduous forest. *American Midland Naturalist* 107:389–392.
- Fellers, J. H. 1987. Interference and exploitation in a guild of woodland ants. *Ecology* 68:1466–1478.
- _____. 1989. Daily and seasonal activity in woodland ants. *Oecologia* 78:69–76.
- Fernández, C. F., and M. L. H. Baena. 1997. Ants of Colombia VII: New species of the genera *Lachnomyrmex* Wheeler and *Megalomyrmex* Forel (Hymenoptera: Formicidae). *Caldasia* 19(1–2): 109–114.
- Fernández, C. F. and E. E. Palacio. 1997. Key to northern South America's *Pogonomyrmex* (Hymenoptera: Formicidae) with description of a new species. *Revista de Biología Tropical* 45(4): 1649–1661.
- Ferreira, L. V., and G. T. Prance. 1998. Species richness and floristic composition in four hectares in the Jaú National Park in upland forests in Central Amazonia. *Biodiversity and Conservation* 7:1349–1364.
- Fisher, B. L. 1996a. Ant diversity patterns along an elevational gradient in the Réserve Naturelle Intégrale d'Andringitra, Madagascar. *Fieldiana: Zoology* (n.s.) 85:93–108.

- _____. 1996b. Origins and affinities of the ant fauna of Madagascar. Pp. 457–465. In W. L. Lourenço (ed.), Biogéographie de Madagascar. Editions ORSTOM, Paris.
- _____. 1997. Biogeography and ecology of the ant fauna of Madagascar (Hymenoptera: Formicidae). *Journal of Natural History* 31:269–302.
- _____. 1998. Ant diversity patterns along an elevational gradient in the Réserve Spéciale d'Anjanaharibe-Sud and on the western Masoala Peninsula, Madagascar. *Fieldiana: Zoology* (n.s.) 90:39–67.
- _____. 1999a. Improving inventory efficiency: A case study of leaf litter diversity in Madagascar. *Ecological Applications* 9:714–731.
- _____. 1999b. Ant diversity patterns along an elevational gradient in the Réserve Naturelle Intégrale d'Andohahela, Madagascar. *Fieldiana: Zoology* (n.s.) 94:129–147.
- Fisher, B. L., and S. Razafimandimbry. 1997. Les fourmis (Hymenoptera: Formicidae). Pp. 104–109. In O. Langrand and S. M. Goodman (eds.), Inventaire Biologique Forêts de Vohibasia et d'Isoky-Vohimena. Recherches pour le Développement, Série Sciences Biologiques No. 12. Centre d'Information et de Documentation Scientifique et Technique and World Wide Fund for Nature, Antananarivo, Madagascar.
- Fisher, B. L., H. Ratsirarson, and S. Razafimandimbry. 1998. Les Fourmis (Hymenoptera: Formicidae). Pp. 107–131. In J. Ratsirarson and S. M. Goodman (eds.), Inventaire Biologique de la Forêt Littorale de Tampolo (Fenoarivo Atsinanana). Recherches pour le Développement, Série Sciences Biologiques No. 14. Centre d'Information et de Documentation Scientifique et Technique and World Wide Fund for Nature. Antananarivo, Madagascar.
- Fittkau, E. J., and H. Klinge. 1973. On biomass and trophic structure of the Central Amazonian rain forest ecosystem. *Biotropica* 5:2–14.
- Fluker, S. S., and J. W. Beardsley. 1970. Sympatric associations of three ants: *Iridomyrmex humilis*, *Pheidole megacephala*, and *Anoplolepis longipes* in Hawaii. *Annals of the Entomological Society of America* 63:1290–1296.
- Forel, A. 1901. Fourmis termitophages, lestobiose, *Atta tardigrada*, sous-genres d'*Euponera*. Annales de la Société Entomologique de Belgique 45:389–398.
- Fowler, H. G. 1988. Taxa of the neotropical grass-cutting ants, *Acromyrmex* (Hymenoptera: Formicidae: Attini). *Científica (Jaboticabal)* 16:281–295.
- _____. 1995. Biodiversity estimates: Ant communities and the rare ant species (Hymenoptera: Formicidae) in a fauna of a sub-tropical island. *Revista de Matemática e Estatística, São Paulo* 13:29–38.
- Fowler, H. G., J. V. E. Bernardi, and L. F. T. di Romagnano. 1990. Community structure and *Solenopsis invicta* in São Paulo. Pp. 199–207. In R. K. Vander Meer, K. Jaffe, and A. Cedeno (eds.), *Applied Myrmecology: A World Perspective*. Westview Press, Boulder, Colorado.
- Fox, M. D., and B. J. Fox. 1982. Evidence for inter-specific competition influencing ant species diversity in a regenerating heathland. Pp. 99–110. In R. C. Buckley (ed.), *Ant-Plant Interactions in Australia*. Dr. W. Junk, The Hague.
- Francoeur, A. 1973. Révision taxonomique des espèces néarctiques du groupe *fusca*, genre *Formica* (Formicidae, Hymenoptera). Mémoires de la Société Entomologique du Québec 3:1–316.
- Francoeur, A. 1985. *Formicoxenus quebecensis* Francoeur sp. nov. Pp. 378–379. In A. Francoeur, R. Loiselle, and A. Buschinger. Biosystématique de la tribu Leptothoracini (Formicidae, Hymenoptera). 1. Le genre *Formicoxenus* dans la région holarctique. *Naturaliste Canadian (Québec)* 112:343–403.
- Franks, N. R., and W. H. Bossert. 1983. The influence of swarm raiding army ants on the patchiness and diversity of a tropical leaf-litter ant community. Pp. 151–163. In S. L. Sutton, T. C. Whitmore, and A. C. Chadwick (eds.), *Tropical Rain Forest: Ecology and Management*. Blackwell, Oxford.
- Friese, C. F., and M. F. Allen. 1988. The interaction of harvester ant activity and VA mycorrhizal fungi. *Proceedings of the Royal Society of Edinburgh* 94B:176.
- _____. 1993. The interaction of harvester ants and vesicular-arbuscular mycorrhizal fungi in a patchy semi-arid environment: The effects of mound structure on fungal dispersion and establishment. *Functional Ecology* 7:13–20.
- Gadagkar, R., K. Chandrashekara, and P. Nair. 1990. Insect species diversity in tropics: Sampling methods and a case study. *Journal of the Bombay Natural History Society* 87(3):337–353.
- Gadagkar, R., P. Nair, K. Chandrashekara, and D. M. Bhat. 1993. Ant species richness and diver-

- sity in some selected localities in Western Ghats, India. *Hexapoda* 5:79–94.
- Gaedike, R., 1995. Colecciones entomologicae (1961–1994). *Nova Supplementa. Entomologica* (Berlin) 6:1–83.
- Gallardo, A. 1929. Note sur les moeurs de la fourmi *Pseudoatta argentina*. *Revista de la Sociedad Entomológica Argentina* 2:197–202.
- Gallardo, J. M. 1951. Sobre um Teiidae (Reptilia, Sauria) poco conocido para la fauna Argentina. *Comunicaciones Instituto Nacional de Investigaciones en Ciencia Naturales* 2:8.
- Gallé, L. 1991. Structure and succession of ant assemblages in a north European sand dune area. *Holarctic Ecology* 14:31–37.
- Gaston, K. 1994. Rarity. Chapman and Hall, New York.
- . 1996. Species richness: Measure and measurement. Pp. 77–113. In K. J. Gaston (ed.), *Biodiversity: A Biology of Numbers and Difference*. Blackwell, Cambridge.
- Gauch, H. G., Jr. 1982. *Multivariate Analysis in Community Structure*. Cambridge University Press, Cambridge.
- Gauld, I., and B. Bolton. 1988. *The Hymenoptera*. Oxford University Press, New York.
- Goeldi, E. 1897. Die Fortpflanzungsweise von 13 brasilianischen Reptilien. *Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere* 10:640–674.
- Goldstein, E. L. 1975. Island biogeography of ants. *Evolution* 29:750–762.
- Goldstein, P. Z. 1999. Functional ecosystems and biodiversity buzzwords. *Conservation Biology* 13:247–255.
- Goncalves, C. R. 1961. O genero *Acromyrmex* no Brasil (Hym. Formicidae). *Studia Entomologica* 4:113–180.
- Gösswald, K. 1932. Ökologische Studien über die Ameisenfauna des mittleren Maingebietes. *Zeitschrift für Wissenschaftliche Zoologie* 142: 1–156.
- Gotelli, N. J. 1993. Ant lion zones: Causes of high-density predator aggregations. *Ecology* 74:226–237.
- . 1996. Ant community structure: Effects of predatory ant lions. *Ecology* 77:630–638.
- Gotwald, W. H., Jr. 1969. Comparative morphological studies of the ants, with particular reference to the mouthparts (Hymenoptera: Formicidae). *Memoirs of the Cornell University Agricultural Experimental Station* 408:1–150.
- . 1982. Army ants. Pp. 157–254. In H. R. Hermann (ed.), *Social Insects*, Vol. 4. Academic Press, New York.
- . 1995. *Army Ants: The Biology of Social Predation*. Cornell University Press, Ithaca, New York.
- Gotwald, W. H., Jr., and W. L. Brown Jr. 1966. The ant genus *Simopelta* (Hymenoptera: Formicidae). *Psyche* (Cambridge) 73:261–277.
- Goulet, H., and J. T. Huber (eds.). 1993. *Hymenoptera of the World: An Identification Guide to Families*. Publication 1894/E. Center for Land and Biological Resources Research, Research Branch, Agriculture Canada, Ottawa, Ontario.
- Greenslade, P., and P. J. M. Greenslade. 1971. The use of baits and preservatives in pitfall traps. *Journal of the Australian Entomological Society* 10:253–260.
- Greenslade, P. J. M. 1964. Pitfall trapping as a method for studying populations of Carabidae (Coleoptera). *Journal of Animal Ecology* 33:301–310.
- . 1971. Interspecific competition and frequency changes among ants in Solomon Islands coconut plantations. *Journal of Applied Ecology* 8:323–352.
- . 1972. Comparative ecology of four tropical ant species. *Insectes Sociaux* 19:195–212.
- . 1973. Sampling ants with pitfall traps: Digging-in effects. *Insectes Sociaux* 20:343–353.
- . 1978. Ants. Pp. 109–113. In W. A. Low (ed.), *The Physical and Biological Features of Kunoth Paddock in Central Australia*. Technical Paper No. 4. CSIRO Division of Land Resources, Canberra.
- . 1979. *A Guide to Ants of South Australia*. South Australian Museum, Adelaide.
- Greenslade, P. J. M., and P. Greenslade. 1977. Some effects of vegetation cover and disturbance on a tropical ant fauna. *Insectes Sociaux* 24:163–182.
- Gregg, R. E. 1959 (1958). Key to the species of *Pheidole* (Hymenoptera: Formicidae) in the United States. *Journal of the New York Entomological Society* 66:7–48.
- . 1963. *The Ants of Colorado*. University of Colorado Press, Boulder.
- Grimaldi, D., Agosti, D., and J. M. Carpenter. 1997. New and rediscovered primitive Ants (Hymenoptera: Formicidae) in Cretaceous Amber from New Jersey, and their phylogenetic Relationships. *American Museum Novitates* 3208:1–43.

- Grime J. P. 1979. Plant Strategies and Vegetation Processes. John Wiley and Sons, Chichester, U.K.
- Haines, B. L. 1978. Element and energy flows through colonies of the leaf-cutting ant, *Atta colombica*, in Panama. *Biotropica* 10:270–277.
- . 1983. Leaf-cutting ants bleed mineral elements out of rainforest in southern Venezuela. *Tropical Ecology* 24:85–93.
- Hamilton, W. D. 1964. The genetical evolution of social behaviour. I. *Journal of Theoretical Biology* 7:1–16.
- Hamilton, W. D. 1972. Altruism and related phenomena, mainly in social insects. *Annual Review of Ecology and Systematics* 3:193–232.
- Handel, S. N., and A. J. Beattie. 1990a. La dispersion des graines par les fourmis. *Pour la Science* 156:54–61.
- . 1990b. Seed dispersal by ants. *Scientific American* 263:76–83.
- Handel, S. N., S. B. Fisch, and G. E. Schatz. 1981. Ants disperse a majority of herbs in a mesic forest community in New York State. *Bulletin of the Torrey Botanical Club* 108:430–437.
- Harada, A. Y., and A. G. Bandeira. 1994. Estratificação e densidade de invertebrados em solo arenoso sob floresta primária e plantios arbóreos na Amazônia central durante estação seca. *Acta Amazonica* 24:103–118.
- Harris, R. A. 1979. A glossary of surface sculpturing. California Department of Food and Agriculture Laboratory Services/Entomology Occasional Papers in Entomology 28:1–31.
- Hashimoto, Y. 1996. Skeletomuscular modifications associated with the formation of an additional petiole on the anterior abdominal segments in aculeate Hymenoptera. *Japanese Journal of Entomology* 64:340–356.
- Hashmi, A. A. 1973. A revision of the Neotropical ant subgenus *Myrmothrix* of genus *Camponotus* (Hymenoptera: Formicidae). *Studia Entomologica* 16:1–140.
- Haskins, C. P., and E. F. Haskins. 1965. *Pheidole megacephala* and *Iridomyrmex humilis* in Bermuda—Equilibrium or slow replacement? *Ecology* 46:736–740.
- . 1988. Final observations on *Pheidole megacephala* and *Iridomyrmex humilis* in Bermuda. *Psyche (Cambridge)* 95:177–184.
- Hayek, L. C., and M. A. Buzas. 1996. Surveying Natural Populations. Columbia University Press, New York.
- Heinze, J., and B. Hölldobler. 1994. Ants in the cold. *Memorabilia Zoologica* 48:99–108.
- Heltshe, J. F., and N. E. Forrester. 1983. Estimating species richness using the jackknife procedure. *Biometrics* 39:1–11.
- Heppner, J. B., and G. Lamas. 1982. Acronyms for world museum collections of insects, with an emphasis on Neotropical Lepidoptera. *Bulletin of the Entomological Society of America* 28: 305–316.
- Heraty, J. M. 1985. A revision of the Nearctic Eucharitinae (Hymenoptera: Chalcidoidea: Eucharitidae). *Proceedings of the Entomological Society of Ontario* 116:61–103.
- . 1986. *Pseudochalcura* (Hymenoptera: Eucharitidae), a New World genus of chalcidoids parasitic on ants. *Systematic Entomology* 11:183–212.
- Heraty, J. M., and D. C. Darling. 1984. Comparative morphology of the planidial larvae of the Eucharitidae and Perilampidae (Hymenoptera: Chalcidoidea). *Systematic Entomology* 9:309–328.
- Herbers, J. M. 1985. Seasonal structuring of a north temperate ant community. *Insectes Sociaux* 32: 224–240.
- . 1989. Community structure in north temperate ants: Temporal and spatial variation. *Oecologia* 81:201–211.
- . 1994. Structure of an Australian ant community with comparisons to North American counterparts (Hymenoptera: Formicidae). *Sociobiology* 24:293–306.
- Herbers, J. M., and S. Grieco. 1994. Population structure of *Leptothorax ambiguus*, a facultatively polygynous and polydomous ant species. *Journal of Evolutionary Biology* 7:581–598.
- Heyer, W. R., M. A. Donnelly, R. W. McDiarmid, L.-A. C. Hayek, and M. S. Foster. 1994. Measuring and Monitoring Biological Diversity: Standard Methods for Amphibians. Smithsonian Institution Press, Washington, D.C.
- Hinkle, G., J. K. Wetterer, T. R. Schultz, and M. L. Sogin. 1994. Phylogeny of the attine ant fungi based on analysis of small subunit ribosomal RNA gene sequences. *Science* 266: 1695–1697.
- Hinton, H. E. 1951. Myrmecophilous Lycaenidae and other Lepidoptera—A summary. *Proceedings and Transactions of the South London Entomological and Natural History Society (1949–1950)*: 111–175.

- Hölldobler, B. 1967. Zur Physiologie der Gast-Wirt-Beziehungen (Myrmecophilie) bei Ameisen. I. Das Gastverhältnis der *Atemeles-* und *Lomechusa*-Larven (Col. Staphylinidae) zu *Formica* (Hym. Formicidae). Zeitschrift für Vergleichende Physiologie 56:1–121.
- . 1968. Verhaltensphysiologische Untersuchungen zur Myrmecophilie einiger Staphylinidenlarven. In W. Herre (ed.), Verhandlungen der Deutschen Zoologischen Gesellschaft (Heidelberg, 1967), Zoologischer Anzeiger, Supplement 31:428–434.
- . 1983. Territorial behavior in the green tree ant (*Oecophylla smaragdina*). Biotropica 15:241–250.
- Hölldobler, B., and C. J. Lumsden. 1980. Territorial strategies in ants. Science 210:732–739.
- Hölldobler, B., and Wilson, E. O. 1977. The number of queens: an important trait in ant evolution. Naturwissenschaften 64:8–15.
- . 1990. The Ants. Belknap Press, Cambridge, Massachusetts.
- Hölldobler, K. 1929. Über eine merkwürdige Parasitenerkrankung von *Solenopsis fugax*. Zeitschrift für Parasitenkunde 2:67–72.
- . 1933. Weitere Mitteilungen über Haplosporidien in Ameisen. Zeitschrift für Parasitenkunde 6:91–100.
- Holway, D. A. 1995. Distribution of the Argentine ant (*Linepithema humile*) in Northern California. Conservation Biology 9:1634–1637.
- Holway, D. A., A. V. Suarez, and T. J. Case. 1998. Loss of intraspecific aggression in the success of a widespread invasive social insect. Science 282:949–952.
- Hood, W. G., and W. R. Tschinkel. 1990. Desiccation resistance in arboreal and terrestrial ants. Physiological Entomology 15:23–35.
- Horn, W., and I. Kahle. 1935a. Über Entomologische Sammlungen, Entomologen, und Entomo-Museologie. Entomologische Beihefte, Berlin-Dahlem.
- . 1935b. Supplement to: Über Entomologische Sammlungen, Entomologen, und Entomo-Museologie. Entomologische Beihefte, Berlin-Dahlem.
- Horn, W., I. Kahle, G. Friese, and R. Gaedike. 1990. Colecciones Entomologicae. Ein Kompendium über den Verbleib Entomologischer Sammlungen der Welt bis 1960. Edition Akademischer Landwirtschaftswissenschaften, Berlin.
- Horvitz, C., and D. Schemske. 1990. Spatiotemporal variation in insect mutualists of a neotropical herb. Ecology 71:1085–1097.
- Hudson, K., and A. Nichols (eds.). 1975. The Directory of World Museums. Columbia University Press, New York.
- Huggert, L., and L. Masner. 1983. A review of myrmecophilic-sympatric diapriid wasps in the Holarctic realm, with descriptions of new taxa and a key to genera (Hymenoptera: Proctotrupoidea: Diapriidae). Contributions of the American Entomological Institute 20:63–89.
- Hughes, R. G. 1986. Theories and models of species abundance. American Naturalist 128:879–899.
- Human, K., and D. Gordon. 1996. Exploitative and interference competition between the Argentine ant and native ant species. Oecologia 105:405–412.
- Hung, A. C. F. 1967. A revision of the ant genus *Polyrhachis* at the subgeneric level. Transactions of the American Entomological Society 93:395–422.
- . 1970. A revision of ants of the subgenus *Polyrhachis* Fr. Smith (Hymenoptera: Formicidae: Formicinae). Oriental Insects 4:1–36.
- Huxley, C. R. 1978. The ant-plants *Myrmecodia* and *Hydnophytum* (Rubiaceae), and the relationships between their morphology, ant occupants, physiology and ecology. New Phytologist 80:231–268.
- Huxley, C. R., and D. F. Cutler (eds.). 1991. Ant-Plant Interactions. Oxford University Press, Oxford.
- Iwanami, Y., and T. Iwadare. 1978. Inhibiting effects of myrmecacin on pollen growth and pollen tube mitosis. Botanical Gazette 139:42–45.
- Jackson, D. A. 1984. Ant distribution patterns in a Cameroonian cocoa plantation: Investigation of the ant mosaic hypothesis. Oecologia 62:318–324.
- Jacobson, E. 1909. Ein Mosquito als Gast und diebischer Schmarotzer der *Cremastogaster diffinis* Smith und eine andere schmarotzende Fliege. Tijdschrift voor Entomologie 52:158–164.
- Jaeger, E. C. 1955. A Source-Book of Biological Names and Terms (3rd ed.). Charles C. Thomas, Springfield, Illinois.
- Janet, C. 1897. Sur les rapports de l'*Antennophorus uhlmanni* Haller avec le *Lasius mixtus* Nyl. Comptes Rendus de l'Académie des Sciences Paris 124:583–585.
- Janos, D. P. 1993. Vesicular-arbuscular mycorrhizae of epiphytes. Mycorrhiza 4:1–4.
- Janzen, D. H. 1966. Coevolution of mutualism between ants and acacias in Central America. Evolution 20:249–275.

- . 1967. Interaction of the bull's-horn acacia (*Acacia cornigera* L.) with an ant inhabitant (*Pseudomyrmex ferruginea* F. Smith) in eastern Mexico. University of Kansas Science Bulletin 47:315–558.
- Janzen, D. H., M. Ataroff, M. Fariñas, S. Reyes, N. Rincon, A. Soler, P. Soriano, and M. Vera. 1976. Changes in the arthropod community along an elevational transect in the Venezuelan Andes. *Biotropica* 8:193–203.
- Jeanne, R. L. 1979. A latitudinal gradient in rates of ant predation. *Ecology* 60:1211–1224.
- Jebb, M. 1991. Cavity structure and function in the tuberous Rubicaceae. Pp. 374–389. In C. R. Huxley and D. F. Cutler (eds.), *Ant-Plant Interactions*. Oxford University Press, Oxford.
- Johnson, R. A. 1992. Soil texture as an influence on the distribution of the desert seed-harvester ants *Pogonomyrmex rugosus* and *Messor pergandei*. *Oecologia* 89:118–124.
- Jolivet, P. 1996. *Ants and Plants: An Example of Coevolution (Enlarged Edition)*. Backhuys, Leiden, Netherlands.
- Jongman, R. H. G., C. J. F. ter Braak, and O. F. R. van Tongeren. 1995. *Data Analysis in Community and Landscape Ecology*. Cambridge University Press, Cambridge.
- Jouvenaz, D. P. 1986. Diseases of fire ants: Problems and opportunities. Pp. 327–338. In C. S. Lofgren and R. K. Vander Meer (eds.), *Fire Ants and Leaf-Cutting Ants: Biology and Management*. Westview Press, Boulder, Colorado.
- Jouvenaz, D. P., and D. W. Anthony. 1979. *Mattesia geminata* sp. n. (Neogregarinida: Ophrocystidae), a parasite of the tropical fire ant, *Solenopsis geminata* (Fabricius). *Journal of Protozoology* 26:354–356.
- Jouvenaz, D. P., G. E. Allen, W. A. Banks, and D. P. Wojcik. 1977. A survey for pathogens of fire ants, *Solenopsis* spp., in the Southeastern United States. *Florida Entomologist* 60:275–279.
- Kane, M. D. 1997. Microbial fermentation in insect guts. Pp. 231–265. In R. I. Mackie and B. A. White (eds.), *Ecology and Physiology of Gastrointestinal Microbes*. Vol. 1. *Gastrointestinal Fermentations and Ecosystems*. Chapman and Hall, New York.
- Karawajew, W. 1906. Weitere Beobachtungen über Arten der Gattung *Antennophorus*. *Mémoires de la Société des Naturalistes de Kiew* 20:209–230.
- Kareiva, P., J. Kingsolver, and R. Huey. 1993. *Biotic Interactions and Global Change*. Sinauer Associates, Sunderland, Massachusetts.
- Kaspari, M. 1993a. Body size and microclimate use in Neotropical granivorous ants. *Oecologia* 96:500–507.
- . 1993b. Removal of seeds from Neotropical frugivore droppings. *Oecologia* 95:81–99.
- . 1996a. Litter ant patchiness at the 1 m² scale: Disturbance dynamics in three Neotropical forests. *Oecologia* 107:265–273.
- . 1996b. Testing resource-based models of patchiness in four Neotropical litter ant assemblages. *Oikos* 76:443–454.
- . 1996c. Worker size and seed size selection by harvester ants in a Neotropical forest. *Oecologia* 105:397–404.
- Kaspari, M., and M. Byrne. 1995. Caste allocation in litter *Pheidole*. *Behavioral Ecology and Sociobiology* 37:255–263.
- Kaspari, M., and E. L. Vargo. 1995. Colony size as a buffer against seasonality: Bergmann's rule in social insects. *American Naturalist* 145:610–632.
- Kaspari, M., and M. Weiser. 1999. The size–grain hypothesis and interspecific scaling in ants. *Functional Ecology* 13:530–538.
- Kaspari, M., and S. Yanoviak. In press. Bait use in tropical litter and canopy ants—evidence for differences in nutrient limitation. *Biotropica*.
- Kaspari, M., L. Alonso, and S. O'Donnell. 2000a. Three energy variables predict ant abundance at a geographic scale. *Proceedings of the Royal Society B* 267:485–490.
- Kaspari, M., S. O'Donnell, and J. Kercher. 2000b. Energy, density, and constraints to species richness: Studies of ant assemblages along a productivity gradient. *American Naturalist* 155:280–293.
- Keeler, K. 1993. Fifteen years of colony dynamics in *Pogonomyrmex occidentalis*, the western harvester ant, in western Nebraska. *Southwestern Naturalist* 38:286–289.
- Kempf, W. W. 1951. A taxonomic study on the ant tribe Cephalotini (Hymenoptera: Formicidae). *Revista Entomologica (Rio de Janeiro)* 22:1–244.
- . 1952. A synopsis of the *pinelii*-complex in the genus *Paracryptocerus* (Hym. Formicidae). *Studia Entomologica* 1:1–30.
- . 1957. Sobre algumas espécies de *Procryptocerus* com a descrição dumha espécie nova (Hymenoptera, Formicidae). *Revista Brasileira de Biologia* 17:395–404.

- _____. 1958a. New studies of the ant tribe Cephalotini (Hym. Formicidae). *Studia Entomologica* (n.s.) 1:1–168.
- _____. 1958b. Estudos sobre *Pseudomyrmex* II. (Hymenoptera: Formicidae). *Studia Entomologica* (n.s.) 1:433–462.
- _____. 1958c. Sobre algumas formigas neotrópicas do gênero *Lepto thorax* Mayr. *Anais da Academia Brasileira de Ciencias* 30:91–102.
- _____. 1959. Two new species of *Gymnomymrmex* Borgmeier, 1954 from southern Brazil, with remarks on the genus (Hymenoptera, Formicidae). *Revista Brasileira de Biologia* 19:337–344.
- _____. 1960a. *Phalacromyrmex*, a new ant genus from southern Brazil (Hymenoptera, Formicidae). *Revista Brasileira de Biologia* 20:89–92.
- _____. 1960b. A review of the ant genus *Mycetarotes* Emery (Hymenoptera, Formicidae). *Revista Brasileira de Biologia* 20:277–283.
- _____. 1960c. Estudo sobre *Pseudomyrmex* I. (Hymenoptera: Formicidae). *Revista Brasileira de Entomologia* 9:5–32.
- _____. 1961a. A survey of the ants of the soil fauna in Surinam (Hymenoptera: Formicidae). *Studia Entomologica* 4:481–524.
- _____. 1961b. Estudos sobre *Pseudomyrmex* III. (Hymenoptera: Formicidae). *Studia Entomologica* 4:369–408.
- _____. 1962. Retoques à classificação das formigas neotropicais do gênero *Heteroponera* Mayr (Hym., Formicidae). *Papéis Avulsos de Zoologia* (São Paulo) 15:29–47.
- _____. 1963. A review of the ant genus *Mycocerpus* Forel, 1893 (Hymenoptera: Formicidae). *Studia Entomologica* 6:417–432.
- _____. 1964. A revision of the Neotropical fungus-growing ants of the genus *Cyphomyrmex* Mayr. Part I: Group of *strigatus* Mayr (Hym., Formicidae). *Studia Entomologica* 7:1–44.
- _____. 1965. A revision of the Neotropical fungus-growing ants of the genus *Cyphomyrmex* Mayr. Part II: Group of *rimosus* (Spinola) (Hym., Formicidae). *Studia Entomologica* 8:161–200.
- _____. 1967a. A synopsis of the Neotropical ants of the genus *Centromyrmex* Mayr (Hymenoptera: Formicidae). *Studia Entomologica* 9:401–410.
- _____. 1967b. Estudos sobre *Pseudomyrmex*. IV (Hymenoptera: Formicidae). *Revista Brasileira de Biologia* 12:1–12.
- _____. 1967c. Three new South American ants (Hym. Formicidae). *Studia Entomologica* 10:353–360.
- _____. 1967d. A new revisionary note on the genus *Paracryptocerus* Emery (Hym. Formicidae). *Studia Entomologica* 10:361–368.
- _____. 1968. A new species of *Cyphomyrmex* from Colombia, with further remarks on the genus (Hymenoptera, Formicidae). *Revista Brasileira de Biologia* 28:35–41.
- _____. 1971. A preliminary review of the ponerine ant genus *Dinoponera* Roger (Hymenoptera: Formicidae). *Studia Entomologica* 14:369–394.
- _____. 1972. Catálogo Abreviado das Formigas da Região Neotropical (Hymenoptera: Formicidae). *Studia Entomologica* 15:3–344.
- _____. 1973a. A revision of the Neotropical myrmicine ant genus *Hylomyrma* Forel (Hymenoptera: Formicidae). *Studia Entomologica* 16:225–260.
- _____. 1973b. A new *Zacryptocerus* from Brazil, with remarks on the generic classification of the tribe Cephalotini (Hymenoptera: Formicidae). *Studia Entomologica* 16:449–462.
- _____. 1974a. A review of the Neotropical ant genus *Oxyepoecus* Santschi (Hymenoptera: Formicidae). *Studia Entomologica* 17:471–512.
- _____. 1975a. A revision of the Neotropical ponerine ant genus *Thaumatomyrmex* Mayr (Hymenoptera: Formicidae). *Studia Entomologica* 18:95–126.
- _____. 1975b. Miscellaneous studies on neotropical ants. VI. (Hymenoptera, Formicidae). *Studia Entomologica* 18:341–380.
- Kempf, W. W., and K. Lenko. 1968. Novas observações e estudos sobre *Gigantiops destructor* (Fabricius) (Hymenoptera: Formicidae). *Papéis Avulsos de Zoologia* (São Paulo) 21:209–230.
- Kent, M., and P. Coker. 1992. *Vegetation Description and Analysis: A Practical Approach*. Belhaven Press, London.
- King, J. R., A. N. Andersen, and A. D. Cutter. 1998. Ants as bioindicators of habitat disturbance: Validation of the functional group model for Australia's humid tropics. *Biodiversity and Conservation* 7:1627–1638.
- Kistner, D. H. 1979. Social and evolutionary significance of social insect symbionts. Pp. 339–413. In H. R. Hermann (ed.), *Social Insects*, Vol. 1. Academic Press, New York.
- _____. 1982. The social insects' bestiary. Pp. 1–244. In H. R. Hermann (ed.), *Social Insects*, Vol. 3. Academic Press, New York.
- Kitching, R. L. 1993. Rainforest canopy arthropods: problems for rapid biodiversity assessment. Pp.

- 26–30. In A. J. Beattie (ed.), *Rapid Biodiversity Assessment: Proceedings of the Biodiversity Assessment Workshop*. Macquarie University, Sydney, Australia.
- Kleinfeldt, S. E. 1978. Ant-gardens: The interaction of *Codonanthe crassifolia* (Gesneriaceae) and *Crematogaster longispina* (Formicidae). *Ecology* 59:449–456.
- . 1986. Ant-gardens: Mutual exploitation. Pp. 283–294. In B. Juniper and T. R. E. Southwood (eds.), *Insects and the Plant Surface*. Edward Arnold, London.
- Koch, C. F. 1987. Prediction of sample size effects on the measured temporal and geographic distribution patterns of species. *Paleobiology* 13:100–107.
- Kohout, R. J. 1988. New nomenclature of the Australian ants of the *Polyrhachis gab* Forel species complex (Hymenoptera: Formicidae: Formicinae). *Australian Entomological Magazine* 15:49–52.
- . 1989. The Australian ants of the *Polyrhachis relucens* species-group (Hymenoptera: Formicidae: Formicinae). *Memoirs of the Queensland Museum* 27:509–516.
- . 1990. A review of the *Polyrhachis viehmeyeri* species-group (Hymenoptera: Formicidae: Formicinae). *Memoirs of the Queensland Museum* 28:499–508.
- Kohout, R. J., and R. W. Taylor. 1990. Notes on Australian ants of the genus *Polyrhachis* Fr. Smith, with a synonymic list of the species (Hymenoptera: Formicidae: Formicinae). *Memoirs of the Queensland Museum* 28:509–522.
- Kremen, C. 1992. Assessing the indicator properties of species assemblages for natural area monitoring. *Ecological Applications* 2:203–217.
- Kremen, C., R. K. Colwell, T. L. Erwin, D. D. Murphy, R. F. Noss, and M. A. Sanjayan. 1994. Terrestrial arthropod assemblages: Their use in conservation planning. *Conservation Biology* 7:796–808.
- Kugler, C. 1978. A comparative study of the myrmicine sting apparatus (Hymenoptera: Formicidae). *Studia Entomologica* 20:413–548.
- Kugler, C. 1994. A revision of the ant genus *Rogeria* with description of the sting apparatus (Hymenoptera: Formicidae). *Journal of Hymenoptera Research* 3:17–89.
- Kugler, C., and W. L. Brown Jr. 1982. Revisionary and other studies on the ant genus *Ectatomma*, including the description of two new species. *Search Agriculture* (Ithaca, N.Y.) 24:1–8.
- Kugler, J. 1986. The Leptanillinae (Hymenoptera: Formicidae) of Israel and a description of a new species from India. *Israel Journal of Entomolgy* 20:45–57.
- Kupyanskaya, A. N. 1980. Ants of the genus *Formica* Linnaeus (Hymenoptera, Formicidae) of the Soviet Far East. Pp. 95–108. In P. A. Ler (ed.), *Taxonomy of Insects of the Far East*. Akademiya Nauk SSSR, Vladivostok. [In Russian.]
- . 1986. Ants (Hymenoptera, Formicidae) of the group *Myrmica lobicornis* Nylander from the Far East. Pp. 83–90. In P. A. Ler (ed.), *Systematics and Ecology of Insects from the Far East*. Akademiya Nauk SSSR, Vladivostok. [In Russian.]
- Kusnezov, N. 1951a. El género *Pogonomyrmex* Mayr (Hym., Formicidae). *Acta Zoologica Lilloana* (Tucuman) 11:227–333.
- . 1951b. *Myrmelachista* en la Patagonia (Hymenoptera, Formicidae). *Acta Zoologica Lilloana* (Tucuman) 11:353–365.
- . 1951c. El género *Pheidole* en la Argentina (Hymenoptera, Formicidae). *Acta Zoologica Lilloana* (Tucuman) 12:5–88.
- . 1951d. El género *Camponotus* en la Argentina (Hymenoptera, Formicidae). *Acta Zoologica Lilloana* (Tucuman) 12:183–252.
- . 1951e. El estado real del grupo *Dorymyrmex* Mayr (Hymenoptera, Formicidae). *Acta Zoologica Lilloana* (Tucuman) 10:427–448.
- . 1955. Zwei neue Ameisengattungen aus Tucuman (Argentinien). *Zoologischer Anzeiger* 154:268–277.
- . 1957. Numbers of species of ants in faunae of different latitudes. *Evolution* 11:298–299.
- Kutter, H. 1931. *Forelophilus*, eine neue Ameisengattung. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 15:193–195.
- . 1945. Ein neue Ameisengattung. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 19:485–487.
- . 1950. Über eine neue, extrem parasitische Ameise. 1. Mitteilung. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 23:81–94.
- . 1973. Zur Taxonomie der Gattung *Chalepoxenus* (Hymenoptera, Formicidae, Myrmicinae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 46:269–280.
- Laakso, J., and H. Setälä. 1997. Nest mounds of red wood ants (*Formica aquilonia*): Hot spots for litter-dwelling earthworms. *Oecologia* 111:565–569.

- Lagerheim, G. 1900. Über *Lasius fuliginosus* und seine Pilzzucht. Entomologisk Tidskrift 21:17–29.
- Lamas, G., R. K. Robbins, and D. J. Harvey. 1991. A preliminary survey of the butterfly fauna of Pakitzá, Parque Nacional del Manu, Peru, with an estimate of its species richness. Publicaciones del Museo de Historia Natural, Universidad Nacional Mayor de San Marcos A 40:1–19.
- Lambbeck, R. J. 1997. Focal species: A multi-species umbrella for nature conservation. Conservation Biology 11:849–856.
- Lambshead, J., and H. M. Platt. 1985. Structural patterns of marine benthic assemblages and their relationships with empirical statistical models. Pp. 371–380. In P. E. Gibbs (ed.), Proceedings of the 19th European Marine Biology Symposium, Plymouth, 1984. Cambridge University Press, Cambridge.
- Landres, P. B., J. Verner, and J. W. Thomas. 1988. Ecological uses of vertebrate indicator species: A critique. Conservation Biology 2:316–328.
- Lattke, J. E. 1986. Two new species of neotropical *Anochetus* Mayr (Hymenoptera: Formicidae). Insectes Sociaux 33:352–358.
- . 1990. A new genus of myrmicine ants (Hymenoptera: Formicidae) from Venezuela. Entomologica Scandinavica 21:173–178.
- . 1991. Studies of neotropical *Amblyopone* Erichson (Hymenoptera: Formicidae). Contributions in Science. Los Angeles County Museum 428:1–7.
- . 1995. Revision of the ant genus *Gnamptogenys* in the New World (Hymenoptera: Formicidae). Journal of Hymenoptera Research 4:137–193.
- . 1997. Revisión del género *Apterostigma* Mayr (Hymenoptera: Formicidae). Arquivos de Zoologia (Museu de Zoologia da Universidade de São Paulo) 34:121–221.
- Lattke, J., and W. Goitia. 1997. El genero *Strumigenys* (Hymenoptera: Formicidae) en Venezuela. Caldasia 19(3):367–396.
- Launer, A. E., and D. D. Murphy. 1994. Umbrella species and the conservation of habitat fragments: A case of a threatened butterfly and a vanishing grassland ecosystem. Biological Conservation 69:145–153.
- Lavorel, S., S. McIntyre, J. Landsberg, and T. D. A. Forbes. 1997. Plant functional classifications: From general groups to specific groups based on response to disturbance. Trends in Ecology and Systematics 12:474–478.
- Lawton, J. 1994. What do species do in ecosystems? Oikos 71:364–374.
- Lawton, J. H., D. E. Bifnell, B. Bolton, G. F. Blommers, P. Eggleton, P. M. Hammond, M. Hodda, R. D. Holt, T. B. Larsen, N. A. Mawdsley, N. E. Stork, D. S. Srivastava, and A. D. Watt. 1998. Biodiversity inventories, indicator taxa and effects of habitat modification in tropical forest. Nature 391:72–76.
- Leary, R., and F. Allendorf. 1989. Fluctuating asymmetry as an indicator of stress: Implications for conservation biology. Trends in Ecology and Evolution 4:214–217.
- Lee, S. M., and Chao, A. 1994. Estimating population size via sample coverage for closed capture-recapture models. Biometrics 50:88–97.
- Le Masne, G. 1941. *Tubicera lichtwardti* Schmitz (Dipt. Phoridae), hôte de *Plagiolepis pygmaea* Latr. (Hym. Formicidae). Bulletin de la Société Entomologique de France 46:110–111.
- Lesica, P. 1993. Using plan community diversity in reserve design for pothole prairie on the Blackfeet Indian Reservation, Montana, USA. Biological Conservation 65:69–75.
- Lesica, P., and P. Kannowski. 1998. Ants create hummocks and alter structure and vegetation of a mountain fen. American Midland Naturalist 139:58–68.
- Lévieux, J. 1976. Étude de la structure du nid de quelques espèces terricoles de fourmis tropicales. Annales de l'Université d'Abidjan, Serie C: Sciences 12:23–33.
- . 1983. The soil fauna of tropical savannas. IV. The ants. Pp. 525–540. In F. Bourlière (ed.), Tropical Savannas: Ecosystems of the World. Elsevier, Amsterdam.
- Levings, S. C. 1983. Seasonal, annual and among-site variation in the ground ant community of a deciduous tropical forest: Some causes of patchy species distributions. Ecological Monographs 53:435–455.
- Levings, S. C., and Traniello, J. F. A. 1981. Territoriality, nest dispersion, and community structure in ants. Psyche (Cambridge) 88:265–319.
- Levy R. 1996. Interspecific colony dispersion and niche relations of three large tropical rain forest ant species. Pp. 331–340. In D. S. Edwards, W. E. Booth, and S. C. Choy (eds.), Tropical Rainforest Research. Kluwer Academic Publishers, Dordrecht.

- Lieberburg, I., P. M. Kranz, and A. Seip. 1975. Bermudian ants revisited: The status and interaction of *Pheidole megacephala* and *Iridomyrmex humilis*. *Ecology* 56:473–478.
- Lincoln, R. J., and G. A. Boxshall. 1987. *The Cambridge Illustrated Dictionary of Natural History*. Cambridge University Press, Cambridge.
- Linnaeus, C. 1758–1759. *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis* (10th ed.). Homiae, Salvii. 2v.
- Littledyke, M., and J. M. Cherrett. 1976. Direct ingestion of plant sap from cut leaves by leaf-cutting ants *Atta cephalotes* (L.) and *Acromyrmex octospinosus* (Reich) (Formicidae, Attini). *Bulletin of Entomological Research* 66:205–217.
- Longino, J. T. 1991. *Azteca* ants in Cecropia trees: Taxonomy, colony structure, and behaviour. Pp. 271–288. In C. R. Huxley and D. F. Cutler (eds.), *Ant-Plant Interactions*. Oxford University Press, Oxford.
- . 1994. How to measure arthropod diversity in a tropical rainforest. *Biology International* 28:3–13.
- Longino, J. T., and R. K. Colwell. 1997. Biodiversity assessment using structured inventory: Capturing the ant fauna of lowland tropical rainforest. *Ecological Applications* 7:1263–1277.
- Longino, J. T., and D. A. Hartley. 1994. *Perissomyrmex snyderi* (Hymenoptera: Formicidae) is native to Central America and exhibits worker polymorphism. *Psyche* (Cambridge) 101:195–202.
- Lubin, Y. D. 1984. Changes in the native fauna of the Galápagos Islands following invasion by the little red fire ant, *Wasmannia auropunctata*. *Biological Journal of the Linnean Society* 21:229–242.
- Ludwig, J. A., and J. F. Reynolds. 1988. *Statistical Ecology: A Primer on Methods and Computing*. John Wiley and Sons, New York.
- Luff, M. L. 1975. Some features influencing the efficiency of pitfall traps. *Oecologia* 19: 345–357.
- Lyford, W. H., 1963. Importance of ants to brown podzolic soil genesis in New England. *Harvard Forest Papers* 7.
- Lynch, J. F. 1981. Seasonal, successional, and vertical segregation in a Maryland ant community. *Oikos* 37:183–198.
- MacArthur, R., and E. O. Wilson. 1967. *The Theory of Island Biogeography*. Princeton University Press, Princeton, New Jersey.
- Mack, A. L. (ed.). 1998. A biological assessment of the Lakekamu basin, Papua New Guinea. RAP Working Papers No. 9. Conservation International, Washington, D.C.
- MacKay, W. P. 1993. A review of the New World ants of the genus *Dolichoderus* (Hymenoptera: Formicidae). *Sociobiology* 22:1–148.
- . 1996. A revision of the ant genus *Acanthostichus* (Hymenoptera: Formicidae). *Sociobiology* 27:129–179.
- . 1997. A revision of the Neotropical ants of the genus *Camponotus*, subgenus *Myrmostenus* (Hymenoptera: Formicidae). *Proceedings of the Entomological Society of Washington* 99(1): 194–203.
- . 2000. A review of the New World ants of the subgenus *Myrafant*, genus *Lepto thorax* (Hymenoptera: Formicidae). *Sociobiology* 36(2): 263–444.
- MacKay, W. P., and E. Mackay. 1997. A revision of the Neotropical ants of the *montivagus* species complex, genus *Camponotus*, subgenus *Myrmecoma* (Hymenoptera: Formicidae). *Sociobiology* 30(3):319–334.
- MacKay, W. P., and S. B. Vinson. 1989. A guide to the species identification of the New World ants. *Sociobiology* 16:3–47.
- Magurran, A. E. 1988. *Ecological Diversity and Its Measurement*. Princeton University Press, Princeton, New Jersey.
- Majer, J. D. 1976. The maintenance of the ant mosaic in Ghana cocoa farms. *Journal of Applied Ecology* 13:123–144.
- . 1980. The influence of ants on broadcast and naturally spread seeds in rehabilitated bauxite mined areas. *Reclamation Review* 3:3–9.
- . 1983. Ants: Bioindicators of minesite rehabilitation, land-use, and land conservation. *Environmental Management* 7(4):375–383.
- . 1984. Recolonisation by ants in rehabilitated open-cut mines in Northern Australia. *Reclamation and Revegetation Research* 2:279–298.
- . 1985. Recolonization by ants of rehabilitated mineral sand mines on North Stradbroke Island, Queensland, with particular reference to seed removal. *Australian Journal of Ecology* 10: 31–48.

- _____. 1990. The abundance and diversity of arboreal ants in northern Australia. *Biotropica* 22: 191–199.
- _____. 1992. Ant recolonization of rehabilitated bauxite mines of Poços de Caldos, Brasil. *Journal of Tropical Ecology* 8:97–108.
- _____. 1996. The use of pitfall traps for sampling ants: A critique. *Proceedings of the Museum of Victoria* 56:323–329.
- Majer, J. D., and A. E. de Kock. 1992. Ant recolonization of sand mines near Richards Bay, South Africa: An evaluation of progress with rehabilitation. *South African Journal of Science* 88: 31–36.
- Majer, J. D., and J. H. C. Delabie. 1994. Comparison of the ant communities of annually inundated and terra firme forests at Trombetas in the Brazilian Amazon. *Insectes Sociaux* 41:343–359.
- Majer, J. D., and O. G. Nichols. 1998. Long-term recolonization patterns of ants in rehabilitated bauxite mines, Western Australia. *Journal of Applied Ecology* 35:161–181.
- Majer, J. D., Day, J. E., Kabay, E. D., and Perriman, W. S. 1984. Recolonization by ants in bauxite mines rehabilitated by a number of different methods. *Journal of Applied Ecology* 21:355–375.
- Majer, J. D., J. H. C. Delabie, and N. L. McKenzie. 1997. Ant litter fauna of forest, forest edges and adjacent grasslands in the Atlantic rain forest region of Bahia, Brazil. *Insectes Sociaux* 44:255–266.
- Malicky, H. 1969. Versuch einer Analyse der ökologischen Beziehungen zwischen Lycaeniden (Lepidoptera) und Formiciden (Hymenoptera). *Tijdschrift voor Entomologie* 112:213–298.
- Mann, W. M. 1916. The Stanford Expedition to Brazil, 1911, John C. Branner, Director: The ants of Brazil. *Bulletin of the Museum of Comparative Zoology, Harvard College* 60:399–490.
- _____. 1921. The ants of the Fiji Islands. *Bulletin of the Museum of Comparative Zoology, Harvard College* 64:401–499.
- _____. 1926. Some new neotropical ants. *Psyche (Cambridge)* 33:97–107.
- Marsh, A. C. 1984. The efficacy of pitfall traps for determining the structure of a desert ant community. *Journal of the Entomological Society of South Africa* 47:115–120.
- _____. 1985. Forager abundance and dietary relationships in a Namib Desert ant community. *South African Journal of Zoology* 20:197–203.
- _____. 1986. Ant species richness along a climatic gradient in the Namib Desert. *Journal of Arid Environments* 11:235–241.
- Martin, J. 1977. Collecting, Preparing, and Preserving Insects, Mites, and Spiders. *The Insects and Arachnids of Canada, Part 1*. Agriculture Canada, Ottawa.
- Maschwitz, U. 1974. Vergleichende Untersuchungen zur Funktion der Ameisenmetathorakaldruse. *Oecologia* 16:303–310.
- Maschwitz, U., and H. Hänel. 1985. The migrating herdsman *Dolichoderus (Diabolus) cuspidatus*: An ant with a novel mode of life. *Behavioral Ecology and Sociobiology* 17:171–184.
- Maschwitz, U., and B. Hölldobler. 1970. Der Kartonnestbau bei *Lasius fuliginosus* Latr. (Hym. Formicidae). *Zeitschrift für Vergleichende Physiologie* 66:176–189.
- Maschwitz, U., and P. Schönegge. 1980. Fliegen als Beute- und Bruträuber bei Ameisen. *Insectes Sociaux* 27:1–4.
- Maschwitz, U., K. Koob, and H. Schildknecht. 1970. Ein Beitrag zur Funktion der Metathorakaldruse der Ameisen. *Journal of Insect Physiology* 16:387–404.
- Masner, L. 1976. Notes on the ectophilous diapriid genus *Mimopria* Holmgren (Hymenoptera: Proctotrupoidea, Diapriidae). *Canadian Entomologist* 108:123–126.
- Masuko, K. 1984. Studies on the predatory biology of oriental Dacetine ants (Hymenoptera: Formicidae). I. Some Japanese species of *Strumigenys*, *Pentastirma*, and *Epitritus*, and a Malaysian *Labidogenys*, with special reference to hunting tactics in short-mandibulate forms. *Insectes Sociaux* 31:429–451.
- _____. 1995 (1994). Specialized predation on orbital mites by two species of the ant genus *Myrmecina* (Hymenoptera: Formicidae). *Psyche (Cambridge)* 101:159–173.
- May, R. M. 1975. Patterns of species abundance and diversity. Pp. 81–120. In M. L. Cody and J. M. Diamond (eds.), *Ecology and Evolution of Communities*. Belknap Press, Cambridge, Massachusetts.
- _____. 1990. How many species? *Philosophical Transactions of the Royal Society of London, Series B* 330 (1257):293–304.
- Mahyé-Nunes, A. 1995. Sinopse do gênero *Mycetarotes* (Hym., Formicidae), com a descrição de duas espécies novas. *Boletim de Entomologia Venezolana* 10(2):197–205.

- Mayr, E. 1942. Systematics and the Origin of Species. Columbia University Press, New York.
- McAreavey, J. 1947. New species of the genera *Prolasius* Forel and *Melophorus* Lubbock (Hymenoptera, Formicidae). Memoirs of the National Museum of Victoria 15:7–27.
- . 1957. Revision of the genus *Stigmacros*. Memoirs of the National Museum of Victoria 21:7–64.
- McArthur, A. J., and M. Adams. 1996. A morphological and molecular revision of the *Camponotus nigriceps* group (Hymenoptera: Formicidae) from Australia. Invertebrate Taxonomy 10(1):1–46.
- McCoy, E. D. 1990. The distribution of insects along elevational gradients. Oikos 58:313–322.
- McGlynn, T. P. 1999a. The biogeography, behavior, and ecology of exotic ants (178 pp.). Ph.D. dissertation, University of Colorado, Boulder, Colorado.
- McGlynn, T. P. 1999b. The worldwide transfer of ants: geographic distribution and ecological invasions. Journal of Biogeography 26:535–548.
- McGlynn, T. P., and C. D. Kelley. 1999. Distribution of a Costa Rican wet forest velvet worm (Onychophora, Peripatidae). Annals of the Entomological Society of America 92:53–55.
- McIlveen, W. D., and H. Cole Jr. 1976. Spore dispersal of Endogonaceae by worms, ants, wasps, and birds. Canadian Journal of Botany 54:1486–1489.
- Medel, R. G., and R. A. Vásquez. 1994. Comparative analysis of harvester ant assemblages of Argentinian and Chilean arid zones. Journal of Arid Environments 26:363–371.
- Menozzi, C. 1929. Revisione delle formiche del genere *Mystrium* Roger. Zoologischer Anzeiger 82:518–536.
- . 1939. Formiche dell'Himalaya e del Karakorum raccolte dalla Spedizione italiana comandata da S. A. R. il Duca di Spoleto (1929). Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale (Milan) 78:285–345.
- Michener, C. D. 1944. Comparative external morphology, phylogeny, and a classification of bees. Bulletin of the American Museum of Natural History 2:151–326.
- Miehe, H. 1911a. Javanische Studien. II. Untersuchungen über die javanische *Myrmecodia*. Abhandlungen der kaiserlichen Sächsischen Gesellschaft der Wissenschaften Mathematisch-physischer Klasse 32:312–361.
- . 1911b. Ueber die javanische *Myrmecodia* und die Beziehung zu ihren Ameisen. Biologisches Zentralblatt 31:733–738.
- Moffett, M. W. 1985. Revision of the genus *Myrmoteras* (Hymenoptera: Formicidae). Bulletin of the Museum of Comparative Zoology, Harvard College 151:1–53.
- . 1986. Revision of the myrmicine genus *Acanthomyrmex* (Hymenoptera: Formicidae). Bulletin of the Museum of Comparative Zoology, Harvard College 151:55–89.
- Moncalvo, J.-M., F. M. Lutzoni, S. A. Rehner, J. Johnson, and R. Vilgalys. 2000. Phylogenetic relationships of agaric fungi based on nuclear large subunit ribosomal DNA sequences. Systematic Biology 49:278–305.
- Morton, S. R., and D. W. Davidson. 1988. Comparative structure of harvester ant communities in arid Australia and North America. Ecological Monographs 58:19–38.
- Moser, J. C. 1964. Inquiline roach responds to trail-marking substance of leaf-cutting ants. Science 143:1048–1049.
- Mueller, U. G., S. A. Rehner, and T. R. Schultz. 1998. The evolution of agriculture in ants. Science 281:2034–2038.
- Munger, J. C. 1992. Reproductive potential of colonies of desert harvester ants (*Pogonomyrmex desertorum*): effects of predation and food. Oecologia 90:276–282.
- Nepstad, D. C., P. Jipp, P. Mautinho, G. Negreiros, and S. Vieira. 1995. Forest recovery following pasture abandonment in Amazonia: Canopy seasonality, fire resistance and ants. Pp. 333–349. In D. J. Rapport, C. L. Gaudet, and P. Calow (eds.): Evaluating and Monitoring the Health of Large-Scale Ecosystems. NATO ASI Series 128.
- New, T. R. 1987. Insect conservation in Australia: Towards rational ecological priorities. Pp. 5–20. In J. D. Majer (ed.), The Role of Invertebrates in Conservation and Biological Survey. Western Australian Department of Conservation and Land Management, Perth, Australia.
- Noss, R. F. 1990. Indicators for monitoring biodiversity: A hierarchical approach. Conservation Biology 4:355–364.
- O'Dowd, D. J. 1982. Pearl bodies as ant food: An ecological role for some leaf emergences of tropical plants. Biotropica 14:40–49.
- Ogata, K. 1982. Taxonomic study of the ant genus *Pheidole* Westwood of Japan, with a description

- of a new species (Hymenoptera, Formicidae). *Kontyû* 50:189–197.
- . 1990. A new species of the ant genus *Epitritus* Emery from Japan (Hymenoptera, Formicidae). *Esakia Special Issue* 1:197–199.
- Ogata, K., and K. Onoyama. 1998. A revision of the ant genus *Smithistruma* Brown of Japan, with descriptions of four new species (Hymenoptera: Formicidae). *Entomological Science* 1(2):277–287.
- Ogata, K., and W. L. Brown Jr. 1991. Ants of the genus *Myrmecia* Fabricius: A preliminary review and key to the named species. *Journal of Natural History* 25:1623–1673.
- Oliveira, P. S. 1988. Ant-mimicry in some Brazilian salticid and clubionid spiders (Araneae: Salticidae, Clubionidae). *Biological Journal of the Linnean Society* 33:1–15.
- Oliveira, P. S., and I. Sazima. 1984. The adaptive bases of ant-mimicry in a Neotropical aphantochilid spider (Araneae: Aphantochilidae). *Biological Journal of the Linnean Society* 22:145–155.
- Oliveira, P. S., M. Galetti, F. Pedroni, and L. P. C. Morellato. 1995. Seed cleaning by *Mycoceropurus goeldii* ants (Attini) facilitates germination in *Hymenaea courbaril* (Caesalpiniaceae). *Biotropica* 27:518–522.
- Oliver, I., and A. J. Beattie. 1996a. Invertebrate morphospecies as surrogates for species: A case study. *Conservation Biology* 10:99–109.
- . 1996b. Designing a cost-effective invertebrate survey: A test of methods for rapid assessment of biodiversity. *Ecological Applications* 6:594–607.
- Oliver I., A. J. Beattie, and A. York. 1998. Spatial fidelity of plant, vertebrate and invertebrate assemblages in multiple-use forest in Eastern Australia. *Conservation Biology* 12(4):822–835.
- Olson, D. M. 1991. A comparison of the efficacy of litter sifting and pitfall traps for sampling leaf litter ants (Hymenoptera: Formicidae) in a tropical wet forest, Costa Rica. *Biotropica* 23:166–172.
- . 1992. Rates of predation by ants (Hymenoptera: Formicidae) in the canopy, understory, leaf litter, and edge habitats of a lowland rainforest in southwestern Cameroon. Pp. 101–109. *In* F. Hall and O. Pascal (eds.), *Biologie d'une Canopie de Forêt Equatoriale II*. Fondation Elf, Paris.
- . 1994. The distribution of leaf litter invertebrates along a Neotropical altitudinal gradient. *Journal of Tropical Ecology* 10:129–150.
- Onoyama, K. 1998. Taxonomic notes on the ant genus *Crematogaster* in Japan (Hymenoptera: Formicidae). *Entomological Science* 1(2):227–232.
- Orlóci, L. 1978. *Multivariate Analysis in Vegetation Research*. Dr. W. Junk, The Hague.
- Orr, A. G., and J. K. Charles. 1994. Foraging in the giant forest ant, *Camponotus gigas* (Smith) (Hymenoptera: Formicidae): Evidence for temporal and spatial specialization in foraging activity. *Journal of Natural History* 28:861–872.
- Orr, M. R. 1992. Parasitic flies (Diptera: Phoridae) influence foraging rhythms and caste division of labor in the leaf-cutter ant *Atta cephalotes* (Hymenoptera: Formicidae). *Behavioral Ecology and Sociobiology* 30:395–402.
- Overal, W. L., and A. G. Bandeira. 1985. Nota sobre hábitos de *Cylindromyrnex striatus* Mayr, 1870, na Amazonia (Formicidae, Ponerinae). *Revista Brasileira de Entomologia* 29:521–522.
- Paine, R. 1968. A note on trophic complexity and community stability. *American Naturalist* 102:91–93.
- Palmer, A., and C. Strobeck. 1986. Fluctuating asymmetry: Measurement, analysis, patterns. *Annual Review of Ecology and Systematics* 17:391–421.
- Palacio, E. 1997. Ants of Colombia VI. Two new species of *Octostruma* (Hymenoptera: Formicidae: Basicerotini). *Caldasia* 19(3):409–418.
- Palmer, M., R. Ambrose, and N. Poff. 1997. Ecological theory and community restoration ecology. *Restoration Ecology* 5:291–300.
- Passera, L. 1994. Characteristics of tramp species. Pp. 23–43. *In* D. F. Williams (ed.), *Exotic Ants: Biology, Impact, and Control of Introduced Species*. Westview Press, Boulder, Colorado.
- Peakall, R., S. N. Handel, and A. J. Beattie. 1991. The evidence for, and importance of, ant pollination. Pp. 421–429. *In* C. R. Huxley and D. F. Cutler (eds.), *Ant-Plant Interactions*. Oxford University Press, Oxford.
- Pearson, D. L., and F. Cassola. 1992. World-wide species richness patterns of tiger beetles (Coleoptera: Cicindelidae): Indicator taxon for biodiversity and conservation studies. *Conservation Biology* 6:376–391.
- Peeters, C. 1991. The occurrence of sexual reproduction among ant workers. *Biological Journal of the Linnean Society* 44:141–152.

- Perfecto, I., and R. R. Snelling. 1995. Biodiversity and the transformation of a tropical agroecosystem: Ants in coffee plantations. *Ecological Applications* 5:1084–1097.
- Petal, J., H. Jakubczyk, A. Breymeyer, and E. Olechowicz. 1971. Productivity investigation of two types of meadows in the Vistula Valley. X. Role of the ants as predators in a habitat. *Ekologia polska* 19:213–222.
- Petersen, B. 1968. Some novelties in presumed males of *Leptanillinae* (Hym., Formicidae). *Entomologiske Meddelelser* 36:577–598.
- Pharo, E. J., A. J. Beattie, and D. Binns. 1999. Vascular plant diversity as a surrogate for bryophyte and lichen diversity. *Conservation Biology* 13:282–292.
- Pielou, E. C. 1966. Species diversity and pattern diversity in the study of ecological succession. *Journal of Theoretical Biology* 10:370–383.
- . 1969. An Introduction to Mathematical Ecology. John Wiley and Sons, New York.
- . 1975. Ecological Diversity. John Wiley and Sons, New York.
- . 1984. The Interpretation of Ecological Data: A Primer on Classification and Ordination. John Wiley and Sons, New York.
- Pierce, N. E. 1987. The evolution and biogeography of association between Lycaenid butterflies and ants. Pp. 89–116. In P. H. Harvey and L. Partridge (eds.), Oxford Surveys in Evolutionary Biology, Vol. 4. Oxford University Press, New York.
- Pierce, N. E., and S. Easteal. 1986. The selective advantage of attendant ants for the larvae of a lycaenid butterfly, *Glaucoopsyche lygdamus*. *Journal of Animal Ecology* 55:451–462.
- Pierce, N. E., and P. S. Mead. 1981. Parasitoids as selective agents in the symbiosis between lycaenid butterfly larvae *Glaucoopsyche lygdamus* and ants. *Outlook* 21:1185–1187.
- Pisarski, B. 1966. Études sur les fourmis du genre *Strongylognathus* Mayr (Hymenoptera, Formicidae). *Annales Zoologici (Warsaw)* 23:509–523.
- Poggi, R., and C. Conci. 1996. Elenco delle collezioni entomologiche conservate nelle strutture pubbliche italiane. *Memorie della Società Entomologica Italiana* 75:3–157.
- Poole, R. W. 1974. An Introduction to Quantitative Ecology. McGraw-Hill, New York.
- Porter, S. D. 1999. FORMIS99: A Master Bibliography of Ant Literature (computer data-base). Published by the author, Gainesville, Florida. <http://cmave.usda.ufl.edu/~formis/>
- Porter, S. D., and M. A. Bowers. 1981. Emigration of an *Atta* colony. *Biotropica* 12:232.
- Porter, S. D., and C. D. Jorgensen. 1988. Longevity of harvester ant colonies in southern Idaho. *Journal of Range Management* 41:104–107.
- Porter, S. D., and D. A. Savignano. 1990. Invasion of polygynous fire ants decimates native ants and disrupts arthropod community. *Ecology* 71:2095–2106.
- Porter, S. D., H. G. Fowler, and W. P. Mackay. 1992. Fire ant mound densities in the United States and Brazil (Hymenoptera: Formicidae). *Journal of Economic Entomology* 85:1154–1161.
- Porter, S. D., H. G. Fowler, S. Campiolo, and M. A. Pesquero. 1995a. Host specificity of several *Pseudacteon* (Diptera: Phoridae) parasites of fire ants (Hymenoptera: Formicidae) in South America. *Florida Entomologist* 78:70–75.
- Porter, S. D., M. A. Pesquero, S. Campiolo, and H. G. Fowler. 1995b. Growth and development of *Pseudacteon* phorid fly maggots (Diptera: Phoridae) in the heads of *Solenopsis* fire ant workers (Hymenoptera: Formicidae). *Environmental Entomology* 24:475–479.
- Porter, S. D., R. K. Vander Meer, M. A. Pesquero, S. Campiolo, and H. G. Fowler. 1995c. *Solenopsis* (Hymenoptera: Formicidae) fire ant reactions to attacks of pseudacteon flies (Diptera: Phoridae) in southeastern Brazil. *Annals of the Entomological Society of America* 88:570–575.
- Prendergast, J. R., R. M. Quinn, J. H. Lawton, B. C. Eversham, and D. W. Gibbons. 1993. Rare species, the coincidence of diversity hotspots and conservation strategies. *Nature* 365:335–337.
- Preston, F. W. 1948. The commonness, and rarity, of species. *Ecology* 29:254–283.
- Prins, A. J. 1982. Review of *Anoplolepis* with reference to male genitalia, and notes on *Acropyga* (Hymenoptera, Formicidae). *Annals of the South African Museum* 89:215–247.
- . 1983. A new ant genus from southern Africa (Hymenoptera, Formicidae). *Annals of the South African Museum* 94:1–11.
- Quinlan, R. J., and J. M. Cherrett. 1979. The role of substrate preparation in the symbiosis between the leaf cutting ant *Acromyrmex octospinosus* (Reich) and its food fungus. *Ecological Entomology* 2:161–170.

- _____. 1979. The role of fungus in the diet of the leaf-cutting ant *Atta cephalotes* (L.). *Ecological Entomology* 4:151–160.
- Quiroz-Robledo, L., and J. Valenzuela-González. 1995. A comparison of ground ant communities in a tropical rainforest and adjacent grassland in Los Tuxtlas, Veracruz, Mexico. *Southwestern Entomologist* 20:203–213.
- Raaijmakers, J. G. W. 1987. Statistical analysis of the Michaelis-Menten equation. *Biometrics* 43:793–803.
- Radchenko, A. G. 1985. Ants of the genus *Strongylognathus* (Hymenoptera: Formicidae) in the European part of the USSR. *Zoologicheskii Zhurnal* 64:1514–1523. [In Russian.]
- _____. 1989a. The ants of the genus *Chalepoxenus* (Hymenoptera, Formicidae) of the USSR fauna. *Vestnik Zoologii* 1989(2):37–41. [In Russian.]
- _____. 1989b. Ants of the *Plagiolepis* genus of the European part of the USSR. *Zoologicheskii Zhurnal* 68(9):153–156. [In Russian.]
- _____. 1991. Ants of the genus *Strongylognathus* (Hymenoptera, Formicidae) of the USSR fauna. *Zoologicheskii Zhurnal* 70(10):84–90. [In Russian.]
- _____. 1992. Ants of the genus *Tetramorium* (Hymenoptera, Formicidae) of the USSR fauna. Report 1. *Zoologicheskii Zhurnal* 71(8):39–49. [In Russian.]
- _____. 1993. Ants of the subfamily Cerapachyinae from Vietnam. *Zhurnal Ukrains'koho Entomolohichnoho Tovarystva* 1:43–47.
- _____. 1994a. Identification table for ants of the genus *Leptothorax* (Hymenoptera, Formicidae) from central and eastern Palearctic. *Zoologicheskii Zhurnal* 73(7–8):146–158. [In Russian.]
- _____. 1994b. A review of the ant genus *Leptothorax* (Hymenoptera, Formicidae) of the central and eastern Palearctic. Communication 1. Subdivision into groups. Groups *acervorum* and *bulgaricus*. *Vestnik Zoologii* 1994(6):22–28. [In Russian.]
- _____. 1995. Palearctic ants of the genus *Cardiomyrmex* (Hymenoptera, Formicidae). *Entomologicheskoe Obozrenie* 74:447–455.
- _____. 1996a. A key of the ant genus *Camponotus* (Hymenoptera, Formicidae) in Palearctic Asia. *Zoologicheskii Zhurnal* 75(8):1195–1203. [In Russian.]
- _____. 1996b. Ants of the genus *Plagiolepis* Mayr (Hymenoptera, Formicidae) of Central and Southern Palaearctic. *Entomologicheskoe Obozrenie* 75(1):178–187.
- _____. 1997. Review of the ants of *scabriceps* group of the genus *Monomorium* Mayr (Hymenoptera, Formicidae). *Annales Zoologici (Warsaw)* 46(3–4):211–224.
- _____. 1998. A key to ants of the genus *Cataglyphis* Foerster (Hymenoptera, Formicidae) of Asia. *Entomologicheskoe Obozrenie* 77(2):502–508, 527. [In Russian].
- Radchenko, A. G., and G. R. Arakelian. 1990. Murav'i gruppy *Tetramorium ferox* Ruzsky iz Kryma i Kavkaza. *Biologicheskii Zhurnal Armenii* 43:371–378.
- Radchenko, A. G., and G. W. Elmes. 1998. Taxonomic revision of the *ritae* species-group of the genus *Myrmica* (Hymenoptera, Formicidae). *Vestnik Zoologii* 32(4):3–27.
- Radchenko, A. G., W. Czechowski, and W. Czechowska. 1997. The genus *Myrmica* Latr. (Hymenoptera, Formicidae) in Poland—A survey of species and a key for their identification. *Annales Zoologici (Warsaw)* 47(3–4): 481–500.
- Rahbek, C. 1995. The elevational gradient of species richness: A uniform pattern? *Ecography* 18:200–205.
- _____. 1997. The relationship among area, elevation, and regional species richness in Neotropical birds. *American Naturalist* 149:875–902.
- Redford, K. H. 1987. Ants and termites as food: Patterns of mammalian myrmecophagy. Pp. 349–399. In H. H. Genoways (ed.), *Current Mammalogy*, Vol. 1. Plenum Press, New York.
- Reichel, H., and A. N. Andersen. 1996. The rainforest ant fauna of Australia's Northern Territory. *Australian Journal of Zoology* 44:81–95.
- Retana, J., X. Cerdá, and X. Espadaler. 1991. Arthropod corpses in a temperate grassland: A limited supply? *Holarctic Ecology* 14:63–67.
- Rettenmeyer, C. W. 1962. The diversity of arthropods found with Neotropical army ants and observations on the behavior of representative species. *Proceedings of the North Central Branch of the Entomological Society of America* 17:14–15.
- _____. 1963. Behavioral studies of army ants. *University of Kansas Science Bulletin* 44:281–465.
- Rettenmeyer, C. W., R. Chadab Crepet, M. G. Naumann, and L. Morales. 1983. Comparative foraging by Neotropical army ants. Pp. 59–73.

- In P. Jaenson (ed.), Social Insects in the Tropics. Université Paris-Nord, Paris.
- Rettig, E. 1904. Ameisenpflanzen-Pflanzenameisen. Beihete zum Botanischen Zentralblatt 17:89–122.
- Rickson, F. R. 1971. Glycogen plastids in Muellerian body cells of *Cecropia peltata*, a higher green plant. Science 173:344–347.
- Rigato, F. 1994a. *Dacatria templaris* gen. n., sp. n. A new myrmicine ant from the Republic of Korea. Deutsche Entomologische Zeitschrift (Neue Folge) 41:155–162.
- _____. 1994b. Revision of the myrmicine ant genus *Lophomyrmex*, with a review of its taxonomic position (Hymenoptera: Formicidae). Systematic Entomology 19:47–60.
- Rissing, S. W. 1987. Annual cycles in worker size of the seed-harvester ant *Veromessor pergandei* (Hymenoptera: Formicidae). Behavioral Ecology and Sociobiology 20:117–124.
- Robertson, H. G. 1990. Unravelling the *Camponotus fulvopilosus* species complex (Hymenoptera: Formicidae). Pp. 327–328. In G. K. Veeresh, B. Mallik, and C. A. Viraktamath (eds.), Social Insects and the Environment. Proceedings of the 11th International Congress of IUSSI, 1990. Oxford and IBH, New Delhi.
- Robinson, S. R., and J. Terborgh. 1990. Bird communities of the Cocha Cashu Biological Station in Amazonian Peru. Frogs, snakes, and lizards of the INPA-WWF reserves near Manaus, Brazil. Pp. 199–216. In A. H. Gentry (ed.), Four Neotropical Rainforests. Yale University Press, New Haven, Connecticut.
- Roepke, W. 1930. Ueber einen merkwürdigen Fall von "Myrmekophilie" bei einer Ameise (*Cladomyrma* sp.?) auf Sumatra beobachtet. Miscellanea Zoologica Sumatrana 45:1–3.
- Rogerson, C. T. 1970. The hypocrealean fungi (Ascomycetes, Hypocreales). Mycologia 62:865–910.
- Rohlfien, K. 1979. Aus der Geschichte der entomologischen Sammlungen des ehemaligen Deutschen Entomologischen Instituts. Beiträge für Entomologie (Berlin) 29:415–438.
- Romero, H., and K. Jaffe. 1989. A comparison of methods for sampling ants (Hymenoptera: Formicidae) in savannas. Biotropica 21: 348–352.
- Room, P. M. 1971. The relative distribution of ant species in Ghana's cocoa farms. Journal of Animal Ecology 40:735–751.
- _____. 1975. Diversity and organization of the ground foraging ant fauna of forest, grassland and tree crops in Papua New Guinea. Australian Journal of Ecology 23:71–89.
- Rosciszewski, K. 1994. *Rostromyrmex*, a new genus of myrmicine ants from peninsular Malaysia (Hymenoptera: Formicidae). Entomologica Scandinavica 25(2):159–168.
- _____. 1995. Die Ameisenfauna eines tropischen Tieflandregenwaldes in Südostasien: Eine faunistisch und ökologische Bestandsaufnahme. Thesis, Johann Wolfgang Goethe Universität, Frankfurt am Main, Germany.
- Rosengren, R., and P. Pamilo. 1983. The evolution of polygyny and polydomy in mound-building *Formica* ants. Acta Entomologica Fennica 24:65–77.
- Rosenzweig, M. L. 1995. Species Diversity in Space and Time. Cambridge University Press, Cambridge.
- Ross, K. G., and J. C. Trager. 1990. Systematics and population genetics of fire ants (*Solenopsis saevissima* complex) from Argentina. Evolution 44: 2113–2134.
- Roth, D. S., I. Perfecto, and B. Rathcke. 1994. The effects of management systems on ground-foraging ant diversity in Costa Rica. Ecological Applications 4:423–436.
- Ryti, R. T., and T. J. Case. 1988a. Field experiments on desert ants: Testing for competition between colonies. Ecology 69:1993–2003.
- _____. 1988b. The regeneration niche of desert ants: Effects of established colonies. Oecologia 75:303–306.
- _____. 1992. The role of neighborhood competition in the spacing and diversity of ant communities. American Naturalist 139:355–374.
- Sabrosky, C. W. 1959. A revision of the genus *Pholeomyia* in North America (Diptera Milichiidae). Annals of the Entomological Society of America 52:316–331.
- Sachtleben, H. 1961. Second supplement to: Über entomologische Sammlungen, Entomologen, und Entomo-Museologie. Beiträge zur Entomologie 11:481–540.
- Sampson, D. A., E. A. Rickart, and P. C. Gonzales. 1997. Ant diversity and abundance along an elevational gradient in the Philippines. Biotropica 29:349–363.
- Santschi, F. 1923a. Revue des fourmis du genre *Brachymyrmex* Mayr. Anales del Museo Nacional de Historia Natural de Buenos Aires 31:650–678.

- _____. 1923b. Descriptions de nouveaux Formicides éthiopiens et notes diverses. I. Revue de Zoologie Africaine 11:259–295.
- _____. 1929. Fourmis du Maroc, d'Algérie et de Tunisie. Bulletin et Annales de la Société Royale Entomologique de Belgique 69:138–165.
- _____. 1936. Fourmis nouvelles ou intéressantes de la République Argentine. Revista Entomología (Rio de Janeiro) 6:402–421.
- _____. 1937. Les sexués du genre *Anillidris* Santschi. Bulletin de la Société Entomologique de France 42:68–70.
- Savolainen, R. 1990. Colony success of the submissive ant *Formica fusca* within territories of the dominant *Formica polyctena*. Ecological Entomology 15:79–85.
- Savolainen, R., and K. Vepsäläinen. 1988. A competition hierarchy among boreal ants: Impact on resource partitioning and community structure. Oikos 51:135–155.
- Schimpfer, A. F. W. 1888. Die Wechselbeziehungen zwischen Pflanzen und Ameisen im tropischen Amerika. Botanische Mitteilungen aus den Tropen (Jena) 1:1–98.
- _____. 1898. Pflanzengeographie auf Physiologischer Grundlage. Pp. 149–170. Jena.
- Schindler, D. 1990. Experimental perturbations of whole lakes as tests of hypotheses concerning ecosystem structure and function. Oikos 57: 25–41.
- Schmid-Hempel, P. 1992. Worker castes and adaptive demography. Journal of Evolutionary Biology 5:1–12.
- Schödl, S. 1998. Taxonomic revision of Oriental *Meranoplus* F. Smith, 1853 (Insecta: Hymenoptera: Formicidae: Myrmicinae). Annalen des Naturhistorischen Museums In Wien. Serie B. Botanik und Zoologie 100B:361–394.
- Schroth, M., and U. Maschwitz. 1984. Zur Larvalbiologie und Wirtsfindung von *Maculinea teleius* (Lepidoptera: Lycaenidae), eines Parasiten von *Myrmica laevinodis* (Hymenoptera: Formicidae). Entomologia Generalis 9:225–230.
- Schulenberg, T. S., and K. Awbrey (eds.). 1997. The Cordillera del Condor of Ecuador and Peru: A biological assessment. RAP Working Papers No. 7. Conservation International, Washington, D.C.
- Schultz, T. R. 1998. Phylogeny of the fungus-growing ants (Myrmicinae: Attini): Evidence from DNA sequences (nuclear elongation factor-1 alpha and mitochondrial cytochrome oxidase I) and morphology. P. 429. In M. P. Schwarz and K. Hogendoorn (eds.), Social Insects at the Turn of the Millennium: Proceedings of the XIII International Congress of IUSSI, Adelaide, Australia, 29 December 1998–3 January 1999. Flinders University Press, Adelaide, Australia.
- Schultz, T. R., and R. Meier. 1995. A phylogenetic analysis of the fungus-growing ants (Hymenoptera: Myrmicinae: Attini) based on morphological characters of the larvae. Systematic Entomology 20:337–370.
- Schumacher, A., and W. G. Whitford. 1976. Spatial and temporal variation in Chihuahuan desert ant faunas. Southwestern Naturalist 21:1–8.
- Schupp, E. W., and D. H. Feener. 1991. Phylogeny, lifeform, and habitat dependence of ant-defended plants in a Panamanian forest. Pp. 175–197. In C. R. Huxley and D. F. Cutler (eds.), Ant-Plant Interactions. Oxford University Press, Oxford.
- Seastedt, T. R., and D. A. Crossley Jr. 1984. The influence of arthropods on ecosystems. BioScience 34:157–161.
- Seifert, B. 1988a. A revision of the European species of the ant subgenus *Chthonolasius* (Insecta, Hymenoptera, Formicidae). Entomologische Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden 51:143–180.
- _____. 1988b. A taxonomic revision of the *Myrmica* species of Europe, Asia Minor, and Caucasia (Hymenoptera, Formicidae). Abhandlungen und Berichte des Naturkundemuseums Görlitz 62(3):1–75.
- _____. 1990. Supplementation to the revision of European species of the ant subgenus *Chthonolasius* Ruzsky, 1913 (Hymenoptera: Formicidae). Doriana 6(271):1–13.
- _____. 1992. A taxonomic revision of the Palaeoarctic members of the ant subgenus *Lasius* s. str. (Hymenoptera: Formicidae). Abhandlungen und Berichte des Naturkundemuseums Görlitz 66(5): 1–67.
- Shattuck, S. O. 1987. An analysis of geographic variation in the *Pogonomyrmex occidentalis* complex (Hymenoptera: Formicidae). Psyche (Cambridge) 94:159–179.
- _____. 1990. Revision of the dolichoderine ant genus *Turneria* (Hymenoptera: Formicidae). Systematic Entomology 15:101–117.
- _____. 1991. Revision of the dolichoderine ant genus *Axinidris* (Hymenoptera: Formicidae). Systematic Entomology 16:105–120.

- _____. 1992a. Review of the dolichoderine ant genus *Iridomyrmex* Mayr with descriptions of three new genera (Hymenoptera: Formicidae). *Journal of the Australian Entomological Society* 31:13–18.
- _____. 1992b. Generic revision of the ant subfamily Dolichoderinae (Hymenoptera: Formicidae). *Sociobiology* 21:1–181.
- _____. 1992c. Higher classification of the ant subfamilies Aneuretinae, Dolichoderinae and Formicinae (Hymenoptera: Formicidae). *Systematic Entomology* 17:199–206.
- _____. 1993. Revision of the *Iridomyrmex purpureus* species-group (Hymenoptera: Formicidae). *Invertebrate Taxonomy* 7:113–149.
- _____. 1994. Taxonomic Catalog of the Ant Subfamilies Aneuretinae and Dolichoderinae (Hymenoptera: Formicidae). University of California Publications in Entomology 112.
- Shattuck, S. O. 1996a. Revision of the *Iridomyrmex discors* species-group (Hymenoptera: Formicidae). *Australian Journal of Entomology* 35(1):37–42.
- Shattuck, S. O. 1996b. The Australian ant genus *Froggattella* Forel (Hymenoptera: Formicidae) revisited. *Australian Journal of Entomology* 35(1):43–47.
- Sheela, S., and T. C. Narendran. 1997. A new genus and a new species of Myrmicinae (Hymenoptera: Formicidae) from India. *Journal of Ecobiology* 9(2):88–91.
- Silva, D., and J. A. Coddington. 1996. Spiders of Pakitza (Madre de Dios, Peru): Species richness and notes on community structure. Pp. 253–311. In D. E. Wilson and A. Sandoval (eds.), *Manu: The Biodiversity of Southeastern Peru*. Smithsonian Institution, Washington, D.C.
- Silvestri, F. 1925 (1924). A new myrmecophilous genus of Coccidae (Hemiptera) from India. *Records of the Indian Museum* 26:311–315.
- Skwarra, E. 1934. Ökologie der Lebensgemeinschaften mexikanischer Ameisenpflanzen. *Zeitschrift für Morphologie und Ökologie der Tiere* 29:306–373.
- Smallwood, J. 1982. Nest relocations in ants. *Insectes Sociaux* 29:138–147.
- Smith, J. B. 1906. An Explanation of Terms Used in Entomology. Brooklyn Entomological Society, Brooklyn, New York.
- Smith, M. R. 1944. The genus *Lachnomyrmex*, with the description of a second species (Hymenoptera: Formicidae). *Proceedings of the Entomological Society of Washington* 46:225–228.
- _____. 1947a. Ants of the genus *Apsychomymex* Wheeler (Hymenoptera: Formicidae). *Revista Entomología (Rio de Janeiro)* 17:468–473.
- _____. 1947b. A new genus and species of ant from Guatemala (Hymenoptera, Formicidae). *Journal of the New York Entomological Society* 55:281–284.
- _____. 1953a. A revision of the genus *Romblonella* W. M. Wheeler (Hymenoptera: Formicidae). *Proceedings of the Hawaiian Entomological Society* 15:75–80.
- _____. 1953b. A new *Romblonella* from Palau, and the first description of a *Romblonella* male (Hymenoptera, Formicidae). *Journal of the New York Entomological Society* 61:163–167.
- _____. 1956a. A key to the workers of *Veromessor* Forel of the United States and the description of a new subspecies (Hymenoptera, Formicidae). *Pacific Entomologist* 32:36–38.
- _____. 1956b. A list of the species of *Romblonella* including two generic transfers (Hymenoptera, Formicidae). *Bulletin of the Brooklyn Entomological Society* 51:18.
- _____. 1961. A study of New Guinea ants of the genus *Aphaenogaster* Mayr (Hymenoptera, Formicidae). *Acta Hymenopterologica* 1:213–238.
- _____. 1962. A remarkable new *Stenamma* from Costa Rica, with pertinent facts on other Mexican and Central American species (Hymenoptera: Formicidae). *Journal of the New York Entomological Society* 70:33–38.
- Smith, T. M., H. H. Shugart, and F. I. Woodward (eds.). 1997. *Plant Functional Types: Their Relevance to Ecosystem Properties and Global Change*. Cambridge University Press, Cambridge.
- Snelling, R. R. 1973. Studies on California ants. 7. The genus *Stenamma* (Hymenoptera: Formicidae). *Contributions in Science (Los Angeles County Museum)* 245:1–38.
- _____. 1975. Descriptions of new Chilean ant taxa (Hymenoptera: Formicidae). *Contributions in Science (Los Angeles County Museum)* 274: 1–19.
- _____. 1976. A Revision of the Honey Ants, Genus *Myrmecocystus* (Hymenoptera: Formicidae). *Los Angeles County Museum of Natural History Bulletin* 24.
- _____. 1979a. Three new species of the Palaeotropical arboreal ant genus *Cataulacus* (Hymenoptera: Formicidae). *Contributions in Science (Los Angeles County Museum)* 274: 1–19.

- noptera: Formicidae). Contributions in Science (Los Angeles County Museum) 315:1–8.
- . 1979b. *Aphomomyrmex* and a related new genus of arboreal African ants (Hymenoptera: Formicidae). Contributions in Science (Los Angeles County Museum) 316:1–8.
- . 1981. The taxonomy and distribution of some North American *Pogonomyrmex* and descriptions of two new species (Hymenoptera: Formicidae). Bulletin of the Southern California Academy of Sciences 80:97–112.
- . 1982. A revision of the honey ants, genus *Myrmecocystus*, first supplement (Hymenoptera: Formicidae). Bulletin of the Southern California Academy of Sciences 81:69–86.
- . 1988. Taxonomic notes on Nearctic species of *Camponotus*, subgenus *Myrmantomia* (Hymenoptera: Formicidae). Pp. 55–78. In J. C. Trager (ed.), Advances in Myrmecology. E. J. Brill, Leiden, Netherlands.
- . 1995a. Systematics of Nearctic ants of the genus *Dorymyrmex* (Hymenoptera: Formicidae). Contributions in Science (Los Angeles County Museum) 454:1–14.
- Snelling, R. R., and J. H. Hunt. 1975. The ants of Chile (Hymenoptera: Formicidae). Revista Chilena de Entomología 9:63–129.
- Snelling, R. R., and J. T. Longino. 1992. Revisionary notes on the fungus-growing ants of the genus *Cyphomyrmex, rimosus* group (Hymenoptera: Formicidae: Attini). Pp. 479–494. In D. Quintero and A. Aiello (eds.), Insects of Panama and Mesoamerica: Selected Studies. Oxford University Press, Oxford.
- Snyder, L. E., and J. M. Herbers. 1991. Polydomy and sexual allocation ratios in the ant *Myrmica punctiventris*. Behavioral Ecology and Sociobiology 28:409–415.
- Soberón, M. J., and J. Llorente B. 1993. The use of species accumulation functions for the prediction of species richness. Conservation Biology 7:480–488.
- Southwood, T. R. E. 1978. Ecological Methods: With Particular Reference to the Study of Insect Populations. Chapman and Hall, London.
- Spellerberg, I. F. 1991. Monitoring Ecological Change. Cambridge University Press, Cambridge.
- . 1992. Evaluation and Assessment for Conservation. Chapman and Hall, London.
- Steghaus-Kovac, S., and U. Maschwitz. 1993. Predation on earwigs: A novel diet specialization within the genus *Leptogenys* (Formicidae, Ponerinae). Insectes Sociaux 40:337–340.
- Stein, M. B., H. G. Thorvilson, and J. W. Johnson. 1990. Seasonal changes in bait preference by the red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae). Florida Entomologist 73: 117–123.
- Stevens, G. C. 1989. The latitudinal gradient in geographical range: How so many species coexist in the tropics. American Naturalist 133:240–256.
- . 1992. The elevational gradient in altitudinal range: An extension of Rapoport's latitudinal rule to altitude. American Naturalist 140:893–911.
- Stork, N. E. 1991. The composition of the arthropod fauna of Bornean lowland rain forest trees. Journal of Tropical Ecology 7:161–180.
- Stork, N. E., and T. M. Blackburn. 1993. Abundance, body size and biomass of arthropods in tropical forest. Oikos 67:483–489.
- Stradling, D. J. 1978. The influence of size on foraging in the ant, *Atta cephalotes*, and the effect of some plant defence mechanisms. Journal of Animal Ecology 47:173–188.
- Sudd, J. H. and N. R. Franks. 1987. The Behavioural Ecology of Ants. Blackwell, Glasgow, U.K.
- Sugihara, G. 1980. Minimal community structure: An explanation of species abundance patterns. American Naturalist 116:770–787.
- Talbot, M. 1943. Population studies of the ant *Prenolepis imparis* Say. Ecology 24:31–44.
- . 1975. A list of the ants of the Edwin George Reserve, Livingston Country, Michigan. Great Lakes Entomologist 8:245–246.
- Taylor, L. R. 1978. Bates, Williams, Hutchinson—A variety of diversities. Pp. 1–18. In L. A. Mound and N. Waloff (eds.), Diversity of Insect Faunas: 9th Symposium of the Royal Entomological Society. Blackwell, Oxford.
- Taylor, L. R., I. P. Woiwod, and J. N. Perry. 1978. The density dependence of spatial behavior and the rarity of randomness. Journal of Animal Ecology 47:383–406.
- Taylor, R. W. 1960. Taxonomic notes on the ants *Ponera leae* Forel and *Ponera norfolkensis* (Wheeler) (Hymenoptera: Formicidae). Pacific Science 14:178–180.
- . 1965. A monographic revision of the rare tropicopolitan ant genus *Probolomyrmex* Mayr (Hymenoptera: Formicidae). Transactions of the Royal Entomological Society of London 117: 345–365.

- _____. 1967. A monographic revision of the ant genus *Ponera* Latreille (Hymenoptera: Formicidae). Pacific Insects Monographs 13:1–112.
- _____. 1968a. Notes on the Indo-Australian basicerotine ants (Hymenoptera: Formicidae). Australian Journal of Zoology 16:333–348.
- _____. 1968b. A new Malayan species of the ant genus *Epitritus*, and a related new genus from Singapore (Hymenoptera: Formicidae). Journal of the Australian Entomological Society 7:130–134.
- _____. 1970a. Characterization of the Australian endemic ant genus *Peronomymex* Viehmeyer (Hymenoptera: Formicidae). Journal of the Australian Entomological Society 9:209–211.
- _____. 1970b. Notes on some Australian and Melanesian basicerotine ants (Hymenoptera: Formicidae). Journal of the Australian Entomological Society 9:49–52.
- _____. 1973. Ants of the Australian genus *Mesostruma* Brown (Hymenoptera: Formicidae). Journal of the Australian Entomological Society 12:24–38.
- _____. 1977. New ants of the Australasian genus *Orectognathus*, with a key to the known species (Hymenoptera: Formicidae). Australian Journal of Zoology 25:581–612.
- _____. 1978a. *Nothomyrmecia macrops*: A living-fossil ant rediscovered. Science 201:979–985.
- _____. 1978b. A taxonomic guide to the ant genus *Orectognathus* (Hymenoptera: Formicidae). CSIRO Division of Entomology Reports 3:1–11.
- _____. 1978c. Melanesian ants of the genus *Amblyopone* (Hymenoptera: Formicidae). Australian Journal of Zoology 26:823–839.
- _____. 1979a. New Australian ants of the genus *Orectognathus*, with summary description of the twenty-nine known species (Hymenoptera: Formicidae). Australian Journal of Zoology 27:773–788.
- _____. 1979b. Notes on the Russian endemic ant genus *Aulacopone* Arnoldi (Hymenoptera: Formicidae). Psyche (Cambridge) 86:353–361.
- _____. 1980. Australian and Melanesian ants of the genus *Eurhopalothrix* Brown and Kempf—Notes and new species (Hymenoptera: Formicidae). Journal of the Australian Entomological Society 19:229–239.
- _____. 1983. Descriptive taxonomy: Past, present and future. Pp. 93–134. In E. Highley and R. W. Taylor (eds.), Australian Systematic Entomology: A Bicentenary Perspective. CSIRO, Melbourne.
- _____. 1985. The ants of the Papuan genus *Dacetinops* (Hymenoptera: Formicidae: Myrmicinae). Series Entomologica (Hague) 33:41–67.
- _____. 1989. Australasian ants of the genus *Leptothorax* Mayr (Hymenoptera: Formicidae: Myrmicinae). Memoirs of the Queensland Museum 27:605–610.
- _____. 1990a. [Untitled. Anomalomyrmini Taylor tribe n., *Anomalomyrma* Taylor gen. n., *Protanilla* Taylor gen. n.] Pp. 278–279. In B. Bolton, The higher classification of the ant subfamily Leptanillinae (Hymenoptera: Formicidae). Systematic Entomology 15:267–282.
- _____. 1990b. New Asian ants of the tribe Basicerotini, with an on-line computer interactive key to the twenty-six known Indo-Australian species (Hymenoptera: Formicidae: Myrmicinae). Invertebrate Taxonomy 4:397–425.
- _____. 1990c. The nomenclature and distribution of some Australian and New Caledonian ants of the genus *Meranoplus* Fr. Smith (Hymenoptera: Formicidae: Myrmicinae). General and Applied Entomology 22:31–40.
- _____. 1990d. Notes on the ant genera *Romblonella* and *Willowsiella*, with comments on their affinities, and the first descriptions of Australian species (Hymenoptera: Formicidae: Myrmicinae). Psyche (Cambridge) 97:281–296.
- Tennant, L. E., and S. D. Porter. 1991. Comparison of diets of two fire ant species (Hymenoptera: Formicidae): Solid and liquid components. Journal of Entomological Science 26:450–465.
- Terayama, M. 1985a. Two new species of the ant genus *Myrmecina* (Insecta; Hymenoptera; Formicidae) from Japan and Taiwan. Edaphologia 32: 35–40.
- _____. 1985b. Two new species of the genus *Acropyga* (Hymenoptera, Formicidae) from Taiwan and Japan. Kontyû 53:284–289.
- _____. 1987. A new species of *Amblyopone* (Hymenoptera, Formicidae) from Japan. Edaphologia 36:31–33.
- Terayama, M. 1996. Taxonomic studies on the Japanese Formicidae, part 2. Seven genera of Ponerinae, Cerapachyinae und Myrmicinae. Nature and Human Activities 1:1–8.
- Terayama, M., and K. Ogata. 1988. Two new species of the ant genus *Probolomyrmex* (Hymenoptera, Formicidae) from Japan. Kontyû 56:590–594.

- Terayama, M. and K. Onoyama. 1999. The ant genus *Leptothorax* Mayr (Hymenoptera: Formicidae) in Japan. Memoirs of the Myrmecological Society of Japan 1:71–97.
- Terayama, M., and S. Yamane. 1989. The army ant genus *Aenictus* (Hymenoptera, Formicidae) from Sumatra, with descriptions of three new species. Japanese Journal of Entomology 57:597–603.
- Terayama, M., C.-C. Lin, and W.-J. Wu. 1995. The ant genera *Epitritus* and *Kyidris* from Taiwan (Hymenoptera: Formicidae). Proceedings of the Japanese Society of Systematic Zoology 53:85–89.
- _____. 1996. The Taiwanese species of the ant genus *Smithistruma* (Hymenoptera, Formicidae). Japanese Journal of Entomology 64(2):327–339.
- Terron, G. 1974. Découverte au Cameroun de deux espèces nouvelles du genre *Prionopelta* Mayr (Hym.: Formicidae). Annales de la Faculté des Sciences, Université Fédéral du Cameroun (Yaoundé) 17:105–119.
- _____. 1981. Deux nouvelles espèces éthiopiennes pour le genre *Proceratium* (Hym.: Formicidae). Annales de la Faculté des Sciences, Université Fédéral du Cameroun (Yaoundé) 28:95–103.
- Thaxter, R. 1888. The Entomophthoreæ of the United States. Memoirs of the Boston Society of Natural History 4:133–201.
- _____. 1908. Contribution toward a monograph of the Laboulbeniaceæ, pt. II. Memoirs of the American Academy of Arts and Sciences 13: 217–469.
- Thompson, C. R., and C. Johnson. 1989. Rediscovered species and revised key to the Florida thief ants (Hymenoptera: Formicidae). Florida Entomologist 72:697–698.
- Tillyard, R. J. 1926. The Insects of Australia and New Zealand. Angus and Robertson, Sydney.
- Tilman, D. 1996. Biodiversity: Population versus ecosystem stability. Ecology 77:350–363.
- Tinaut, A. 1990. *Teleutomyrmex kutteri*, spec. nov.: A new species from Sierra Nevada (Granada, Spain). Spixiana 13:201–208.
- Tobin, J. E. 1991. A neotropical rain forest canopy ant community: Some ecological considerations. Pp. 536–538. In C. R. Huxley and D. F. Cutler (eds.), Ant-Plant Interactions. Oxford University Press, Oxford.
- _____. 1994. Ants as primary consumers: Diet and abundance in the Formicidae. Pp 279–307. In J. H. Hunt and C. A. Nalepa (eds.), Nourishment and Evolution in Insect Societies. Westview Press, Boulder, Colorado.
- _____. 1997. Competition and coexistence of ants in a small patch of rainforest canopy in Peruvian Amazonia. Journal of the New York Entomological Society 105:105–112.
- Tohmé, G., and H. Tohmé. 1981. Les fourmis du genre *Messor* en Syrie. Position systématique. Description de quelques ailés et de formes nouvelles. Répartition géographique. Ecologia Mediterranea 7(1):139–153.
- Topoff, H. 1990. Slave-making ants. American Scientist 78:520–528.
- Torre-Bueno, J. R. de la. 1937. A Glossary of Entomology. Brooklyn Entomological Society, Brooklyn, New York.
- Torre-Bueno, J. R. de la. 1989. The Torre-Bueno Glossary of Entomology. Compiled by S.W. Nichols, and including Supplement A by G. S. Tulloch. New York Entomological Society, New York.
- Trager, J. C. 1984. A revision of the genus *Paratrechina* (Hymenoptera: Formicidae) of the continental United States. Sociobiology 9:49–162.
- _____. 1991. A revision of the fire ants, *Solenopsis geminata* group (Hymenoptera: Formicidae: Myrmicinae). Journal of the New York Entomological Society 99:141–198.
- Trivers, R. L., and H. Hare. 1976. Haplodiploidy and the evolution of the social insects. Science 191:249–263.
- Tschinkel, W. R. 1991. Insect sociometry, a field in search of data. Insectes Sociaux 38:77–82.
- _____. 1992. Brood raiding and the population dynamics of founding and incipient colonies of the fire ant, *Solenopsis invicta*. Ecological Entomology 17:179–188.
- _____. 1993. Sociometry and sociogenesis of colonies of the fire ant *Solenopsis invicta* during one annual cycle. Ecological Monographs 63: 425–457.
- Tschinkel, W. R., and Howard, D. F. 1978. Queen replacement in orphaned colonies of the fire ant, *Solenopsis invicta*. Behavioral Ecology and Sociobiology 3:297–310.
- Tulloch, G. S. 1962. Torre-Bueno's Glossary of Entomology, Supplement A. Brooklyn Entomological Society, Brooklyn, New York.
- Turk, F. A. 1953. A new genus and species of pseudoscorpion with some notes on its biology. Proceedings of the Zoological Society of London 122:951–954.

- Ule, E. 1902. Ameisengärten im Amazonasgebiet. *Botanische Jahrbücher für Systematik, Pflanzen-geschichte und Pflanzengeographien* 30:45–52.
- Umphrey, G. J., 1996. Morphometric discrimination among sibling species in the *fulva-rudis-texana* complex of the ant genus *Aphaenogaster* (Hymenoptera: Formicidae). *Canadian Journal of Zoology* 74(3):528–559.
- United Nations Environment Programme. 1995. *Global Biodiversity Assessment*. Cambridge University Press, Cambridge.
- Upton, M. 1991. Methods for Collecting, Preserving and Studying Insects and Allied Forms. Australian Entomological Society Miscellaneous Publications [Brisbane, Australia] 3:1–86.
- Valone, T., and J. Brown. 1995. Effects of competition, colonization, and extinction on rodent species diversity. *Science* 267:880–883.
- Vander Meer, R., and L. Alonso. 1998. Pheromone directed behavior in ants. Pp. 159–192. In R. Vander Meer, M. Breed, M. Winston, and K. Espelie (eds.), *Pheromone Communication in Social Insects*. Westview Press, Boulder, Colorado.
- Vasconcelos, H. L., and J. H. C. Delabie. 2000. Ground ant communities from central Amazonia forest fragments. Pp. 59–70. In D. Agosti, J. Majer, L. Alonso, and T. R. Schultz (eds.), *Sampling Ground-Dwelling Ants: Case Studies from the Worlds' Rain Forests*. Curtin University School of Environmental Biology Bulletin No. 18. Perth, Australia.
- Vaz-Ferreira, R., L. C. de Zolessi, and F. Achával. 1970. Oviposicion y desarrollo de Ofidios y Lacertilios en hormigueros de *Acromyrmex*. *Physis* 29:431–459.
- . 1973. Oviposición y desarrollo de ofidos y lacertilios en hormigueros de *Acromyrmex*. II. *Trabajos del Cinco Congresso Latinoamericano de Zoología*, Montevideo 1:232–244.
- Veerhagh, M. 1990. The Formicidae of the rain forest in Panguana, Peru: The most diverse local ant fauna ever recorded. Pp. 217–218. In G. K. Veeresh, B. Mallik, and C. A. Viraktamath (eds.), *Social Insects and the Environment*. Proceedings of the 11th International Congress of IUSSI, 1990. Oxford and IBH, New Delhi.
- Vinson, S. B. 1991. Effect of the red imported fire ant (Hymenoptera: Formicidae) on a small plant-decomposing arthropod community. *Environmental Entomology* 20:98–103.
- Von Ihering, H. 1891. Die Wechselbeziehungen zwischen Pflanzen und Ameisen in den Tropen. *Ausland* 1891:474–477.
- Wang, C., and J. Wu. 1991. Taxonomic studies on the genus *Polyrhachis* Mayr of China (Hymenoptera, Formicidae). *Forest Research* 4:596–601. [In Chinese.]
- Wang, C., G. Xiao, and J. Wu. 1989a. Taxonomic studies on the genus *Camponotus* Mayr in China (Hymenoptera, Formicidae) [part]. *Forest Research* 2:221–228. [In Chinese.]
- . 1989b. Taxonomic studies on the genus *Camponotus* Mayr in China (Hymenoptera, Formicidae). [conclusion]. *Forest Research* 2:321–328. [In Chinese.]
- Wang, M. 1993. Taxonomic study of the ant tribe Odontomachini in China (Hymenoptera: Formicidae). *Scientific Treatise on Systematic and Evolutionary Zoology* 2:219–230. [In Chinese.]
- Wang, M., G. Xiao, and J. Wu. 1988. Taxonomic studies on the genus *Tetramorium* Mayr in China (Hymenoptera, Formicidae). *Forest Research* 1:264–274. [In Chinese.]
- Ward, P. S. 1980. A systematic revision of the *Rhytidoponera impressa* group (Hymenoptera: Formicidae) in Australia and New Guinea. *Australian Journal of Zoology* 28:475–498.
- . 1984. A revision of the ant genus *Rhytidoponera* (Hymenoptera: Formicidae) in New Caledonia. *Australian Journal of Zoology* 32:131–175.
- . 1985. The Nearctic species of the genus *Pseudomyrmex* (Hymenoptera: Formicidae). *Quaestiones Entomologicae* 21:209–246.
- . 1987. Distribution of the introduced Argentine ant (*Iridomyrmex humilis*) in natural habitats of the lower Sacramento Valley and its effects on the indigenous ant fauna. *Hilgardia* 55:1–16.
- . 1988. Mesic elements in the western Nearctic ant fauna: Taxonomic and biological notes on *Amblyopone*, *Proceratium*, and *Smithistruma* (Hymenoptera: Formicidae). *Journal of the Kansas Entomological Society* 61:102–124.
- . 1989. Systematic studies on pseudomyrmecine ants: Revision of the *Pseudomyrmex ocularis* and *P. subtilissimus* species groups, with taxonomic comments on other species. *Quaestiones Entomologicae* 25:393–468.
- . 1990. The ant subfamily Pseudomyrmicinae (Hymenoptera: Formicidae): Generic revi-

- sion and relationship to other formicids. *Systematic Entomology* 15:449–489.
- _____. 1993. Systematic studies on *Pseudomyrmex* acacia-ants (Hymenoptera: Formicidae: Pseudomyrmecinae). *Journal of Hymenoptera Research* 2:117–168.
- _____. 1994. *Adetomyrma*, an enigmatic new ant genus from Madagascar (Hymenoptera: Formicidae), and its implications for ant phylogeny. *Systematic Entomology* 19:159–175.
- _____. 1999a. Systematics, biogeography and host plant associations of the *Pseudomyrmex vidiuus* group (Hymenoptera: Formicidae), *Triplaris-* and *Tachigali*-inhabiting ants. *Zoological Journal of the Linnean Society* 126:451–540.
- _____. 1999b. Deceptive similarity in army ants of the genus *Neivamyrmex* (Hymenoptera: Formicidae): Taxonomy, distribution and biology of *N. californicus* (Mayr) and *N. nigrescens* (Cresson). *Journal of Hymenoptera Research* 8:74–97.
- Ward, P. S., B. Bolton, S. O. Shattuck, and W. L. Brown Jr. 1996. A Bibliography of Ant Systematics. University of California Publications in Entomology 116.
- Wasemann, E. 1902. Zur Kenntnis der myrmecophilen *Antennophorus* und anderer auf Ameisen und Termiten reitende Acarinen. *Zoologischer Anzeiger* 25:66–76.
- Watkins, J. F. II. 1976. The identification and distribution of New World army ants (Dorylinae: Formicidae). Baylor University Press, Waco, Texas.
- _____. 1977. The species and subspecies of *Nomamyrmex* (Dorylinae: Formicidae). *Journal of the Kansas Entomological Society* 50:203–214.
- _____. 1982. The army ants of Mexico (Hymenoptera: Formicidae: Ecitoninae). *Journal of the Kansas Entomological Society* 55:197–247.
- _____. 1985. The identification and distribution of the army ants of the United States of America (Hymenoptera, Formicidae, Ecitoninae). *Journal of the Kansas Entomological Society* 58:479–502.
- Watt, J. C. 1979. Abbreviations for entomological collections. *New Zealand Journal of Zoology* 6:519–520.
- Weber, N. A. 1943. Parabiosis in Neotropical “ant gardens.” *Ecology* 24:400–404.
- _____. 1944. The neotropical coccid-tending ants of the genus *Acropyga* Roger. *Annals of the Entomological Society of America* 37:89–122.
- _____. 1947. A revision of the North American ants of the genus *Myrmica* Latreille with a synopsis of the Palearctic species. I. *Annals of the Entomological Society of America* 40:437–474.
- _____. 1948. A revision of the North American ants of the genus *Myrmica* Latreille with a synopsis of the Palearctic species. II. *Annals of the Entomological Society of America* 41:267–308.
- _____. 1950a. The African species of the genus *Oligomyrmex* Mayr (Hymenoptera, Formicidae). *American Museum Novitates* 1442:1–19.
- _____. 1950b. A revision of the North American ants of the genus *Myrmica* Latreille with a synopsis of the Palearctic species. III. *Annals of the Entomological Society of America* 43:189–226.
- _____. 1952. Studies on African Myrmicinae. I (Hymenoptera, Formicidae). *American Museum Novitates* 1548:1–32.
- _____. 1972a. The Attines: The fungus-culturing ants. *American Scientist* 60:448–456.
- _____. 1972b. Gardening Ants: The Attines. American Philosophical Society, Philadelphia.
- Weber, N. A., and J. L. Anderson. 1950. Studies on central African ants of the genus *Pseudolasius* Emery (Hymenoptera, Formicidae). *American Museum Novitates* 1443:1–7.
- Wehner, R., A. C. Marsh, and S. Wehner. 1992. Desert ants on a thermal tightrope. *Nature* 357:586–587.
- Went, F. W., J. Wheeler, and G. C. Wheeler. 1972. Feeding and digestion in some ants (*Veromessor* and *Manica*). *BioScience* 22:82–88.
- Westman, W. 1986. Resilience: Concepts and measures. Pp. 5–19. In B. Dell, A. Hopkins, and B. Lamont (eds.), *Resilience in Mediterranean Ecosystems*. Dr. W. Junk, The Hague.
- Wetterer, J. K. 1991. Allometry and the geometry of leaf-cutting in *Atta cephalotes*. *Behavioral Ecology and Sociobiology* 29:347–351.
- Wetterer, J. K., T. R. Schultz, and R. Meier. 1998. Phylogeny of fungus-growing ants (tribe Attini) based on mtDNA sequence and morphology. *Molecular Phylogenetics and Evolution* 9: 42–47.
- Wheeler, D., and S. Levings. 1988. The impact of the 1983 El Niño drought on the litter arthropods of Barro Colorado Island, Panama. Pp. 309–326. In J. C. Trager (ed.), *Advances in Myrmecology*. E. J. Brill, New York.
- Wheeler, G. C., and E. W. Wheeler. 1930. Two new ants from Java. *Psyche* (Cambridge) 37:193–201.

- Wheeler, G. C., and J. Wheeler. 1986. The Ants of Nevada. Natural History Museum of Los Angeles County, Los Angeles.
- Wheeler, J. 1968. Male genitalia and the taxonomy of *Polyergus* (Hymenoptera: Formicidae). Proceedings of the Entomological Society of Washington 70:156–164. [Erratum: Proceedings of the Entomological Society of Washington 70:254.]
- Wheeler, W. M. 1905. The North American ants of the genus *Liometopum*. Bulletin of the American Museum of Natural History 21:321–333.
- . 1908. Studies on myrmecophiles. II. *Hetaerius*. Journal of the New York Entomological Society 16:135–143.
- . 1910. Two new myrmecophilous mites of the genus *Antennophorus*. Psyche (Cambridge) 17:1–6.
- . 1913. Observations on the Central American *Acacia* ants. Transactions of the Second International Entomological Congress of Oxford (1912) 2:109–139.
- . 1914. Notes on the habits of *Liomyrmex*. Psyche (Cambridge) 21:75–76.
- . 1918. The Australian ants of the ponerine tribe Cerapachyini. Proceedings of the American Academy of Arts and Sciences 53:215–265.
- . 1922a. Ants of the American Museum Congo expedition. A contribution to the myrmecology of Africa. II. The ants collected by the American Museum Congo Expedition. Bulletin of the American Museum of Natural History 45:39–269.
- . 1922b. Ants of the American Museum Congo expedition. A contribution to the myrmecology of Africa. VII. Keys to the genera and subgenera of ants. Bulletin of the American Museum of Natural History 45:631–710.
- . 1924. Ants of Krakatau and other islands in the Sunda Strait. Treubia 5:239–258.
- . 1925. A new guest-ant and other new Formicidae from Barro Colorado Island, Panama. Biological Bulletin of the Marine Biological Laboratory (Woods Hole) 49:150–181.
- . 1928. The Social Insects: Their Origin and Evolution. Kegan Paul, Trench, Trubner, London.
- . 1934. A second revision of the ants of the genus *Leptomyrmex* Mayr. Bulletin of the Museum of Comparative Zoology, Harvard College 77:69–118.
- . 1935. Ants of the genus *Acropyga* Roger, with description of a new species. Journal of the New York Entomological Society 43:321–329.
- . 1936. Ecological relations of ponerine and other ants to termites. Proceedings of the American Academy of Arts and Sciences 71:159–243.
- . 1942. Studies of Neotropical ant-plants and their ants. Bulletin of the Museum of Comparative Zoology, Harvard University 90:1–262.
- Wheeler, W. M. and W. M. Mann. 1942. [Untitled. *Allomerus decemarticulatus* Mayr subsp. *Nove-marticulatus* Wheeler & Mann, subsp. nov.] Pp. 188–189. In W. M. Wheeler, Studies of Neotropical ant-plants and their ants. Bulletin of the Museum of Comparative Zoology, Harvard College 90:1–262.
- Whitcomb, W. H., A. Bhatkar, and J. C. Nickerson. 1973. Predators of *Solenopsis invicta* queens prior to successful colony establishment. Environmental Entomology 2:1101–1103.
- Whitford, W. G. 1978. Structure and seasonal activity of Chihuahua desert ant communities. Insectes Sociaux 25:79–88.
- Whitford, W. G., and G. Ettershank. 1975. Factors affecting foraging activity in Chihuahuan desert harvester ants. Environmental Entomology 4: 689–696.
- Wiernasz, D. C., and B. J. Cole. 1995. Spatial distribution of *Pogonomyrmex occidentalis*: Recruitment, mortality, and overdispersion. Journal of Animal Ecology 64:519–527.
- Wilcox, B. A. 1984. In situ conservation of genetic resources: Determinants of minimum-area requirements. Pp. 639–647. In J. A. McNeely and K. R. Miller (eds.), National Parks, Conservation, and Development: The Role of Protected Areas in Sustaining Society. Proceedings of the World Congress on National Parks, Bali, Indonesia, 11–22 October 1982. Smithsonian Institution Press, Washington, D.C.
- Wilcox, B. A., D. D. Murphy, P. R. Ehrlich, and G. T. Austin. 1986. Insular biogeography of the montane butterfly faunas in the Great Basin: Comparison with birds and mammals. Oecologia 69:188–194.
- Willey, R. B., and W. L. Brown Jr. 1983. New species of the ant genus *Myopias* (Hymenoptera: Formicidae: Ponerinae). Psyche (Cambridge) 90:249–285.
- Williams, D. F. 1994. Exotic ants: Biology, impact, and control of introduced species. Westview Press, Boulder, Colorado.
- Williams, R. N. (ed.). 1978. Worldwide directory of institutions with entomologists, part I: Latin

- America. Bulletin of the Entomological Society of America 24:179–193.
- Willis, E. O. 1983. A study of ant-following birds of northeastern Brazil. Research Reports of the National Geographic Society 15:745–748.
- Willis, E. O., and Y. Oniki. 1978. Birds and army ants. Annual Review of Ecology and Systematics 9:243–263.
- Wilson, D. E., F. R. Cole, J. D. Nichols, R. Rudran, and M. S. Foster. 1996. Measuring and Monitoring Biological Diversity. Standard Methods for Mammals. Smithsonian Institution Press, Washington, D.C.
- Wilson, E. O. 1953. The ecology of some North American dacetine ants. Annals of the Entomological Society of America 46:479–495.
- . 1955. A monographic revision of the ant genus *Lasius*. Bulletin of the Museum of Comparative Zoology, Harvard College 113:1–201.
- . 1958. Patchy distributions of ant species in New Guinea rain forests. Psyche (Cambridge) 65:26–38.
- . 1959. Some ecological characteristics of ants in New Guinea rain forests. Ecology 40:437–447.
- . 1961. The nature of the taxon cycle in the Melanesian ant fauna. American Naturalist 95: 169–193.
- . 1962a. Behavior of *Dacetum armigerum* (Latreille), with a classification of self-grooming movements in ants. Bulletin of the Museum of Comparative Zoology, Harvard College 127:401–422.
- . 1962b. The Trinidad cave ant *Erebomyrma* (= *Spelaeomyrmex*) *urichi* (Wheeler), with a comment on cavernicolous ants in general. Psyche (Cambridge) 69:63–72.
- . 1964. The true army ants of the Indo-Australian area (Hymenoptera: Formicidae: Dorylinae). Pacific Insects Monographs 6:427–483.
- . 1971. The Insect Societies. Belknap Press, Cambridge, Massachusetts.
- . 1976. Which are the most prevalent ant genera? Studia Entomologica 19:187–200.
- . 1984. Tropical social parasites in the ant genus *Pheidole*, with an analysis of the anatomical parasitic syndrome (Hymenoptera: Formicidae). Insectes Sociaux 31:316–334.
- . 1985. The principles of caste evolution. Fortschritte der Zoologie 31:307–324.
- . 1987. The arboreal ant fauna of Peruvian Amazon forests: A first assessment. Biotropica 19:245–251.
- . 1989. *Chimaeridris*, a new genus of hook-mandibled myrmicine ants from tropical Asia (Hymenoptera: Formicidae). Insectes Sociaux 36:62–69.
- . 1993. The Diversity of Life. W.W. Norton, New York.
- Wilson, E. O., and W. L. Brown Jr. 1953. The subspecies concept and its taxonomic application. Systematic Zoology 2:97–111.
- . 1956. New parasitic ants of the genus *Kyidris*, with notes on ecology and behavior. Insectes Sociaux 3:439–454.
- . 1984. Behavior of the cryptobiotic predaceous ant *Eurhopalothrix heliscata* n. sp. (Hymenoptera: Formicidae: Basicerotini). Insectes Sociaux 31:408–428.
- Wilson, E. O., and R. W. Taylor. 1967. An estimate of the potential evolutionary increase in species density in the Polynesian ant fauna. Evolution 21:1–10.
- Wilson, E. O., T. Eisner, G. C. Wheeler, and J. Wheeler. 1956. *Aneuretus simoni* Emery, a major link in ant evolution. Bulletin of the Museum of Comparative Zoology, Harvard College 115:81–99.
- Wing, M. W. 1968. Taxonomic revision of the Nearctic genus *Acanthomyops* (Hymenoptera: Formicidae). Memoirs of the Cornell University Agricultural Experiment Station 405:1–173.
- Wisdom, W., and W. G. Whitford. 1981. Effects of vegetation change on ant communities of arid rangelands. Environmental Entomology 10:893–897.
- Wolda, H. 1992. Trends in abundance of tropical forest insects. Oecologia 89:47–52.
- Wu, J. 1990. Taxonomic studies on the genus *Formica* L. of China (Hymenoptera: Formicidae). Forest Research 3:1–8. [In Chinese.]
- Wu, J., and C. Wang. 1990. A taxonomic study on the genus *Tetraponera* Smith in China (Hymenoptera: Formicidae). Scientia Silvae Sinica 26: 515–518. [In Chinese.]
- Xu, Z. 1994a. A taxonomic study of the ant genus *Lepisiota* Santschi from Southwestern China (Hymenoptera: Formicidae: Formicinae). Journal of Southwest Forestry College 14(4):231–237.
- Xu, Z. 1994b. A taxonomic study of the ant genus *Brachyponera* Emery in Southwestern China (Hymenoptera: Formicidae: Ponerinae). Journal of Southwest Forestry College 14(3):181–185.
- Xu, Z. 1995a. Two new species of the ant genus *Prenolepis* from Yunnan China (Hymenoptera: Formicidae). Zoological Research 16(4):337–341.

- Xu, Z. 1995b. A taxonomic study of the ant genus *Dolichoderus* Lund in China (Hymenoptera: Formicidae: Dolichoderinae). *Journal of Southwest Forestry College* 15(1):33–39.
- Xu, Z. 1997. A taxonomic study of the ant genus *Pseudolasius* Emery in China (Hymenoptera: Formicidae). *Zoological Research* 18(1):1–6.
- Xu, Z. 1999. Systematic studies on the ant genera of *Carebara*, *Rhopalomastix* and *Kartidris* in China (Hymenoptera: Formicidae: Myrmicinae). *Acta Biologica Plateau Sinica* 14:129–136.
- Xu, Z., and W. Zhang. 1996. A new species of the genus *Gnamptogenys* (Hymenoptera: Formicidae: Ponerinae) from southwestern China. *Entomotaxonomia* 18(1):55–58.
- Yamauchi, K. 1978. Taxonomical and ecological studies on the ant genus *Lasius* in Japan (Hymenoptera: Formicidae). I. Taxonomy. *Science Reports of the Faculty of Education, Gifu University (Natural Sciences)* 6:147–181.
- Yasumatsu, K., and W. L. Brown Jr. 1951. Revisional notes on *Camponotus herculeanus* Linné and close relatives in Palearctic regions (Hymenoptera: Formicidae). *Journal of the Faculty of Agriculture, Kyushu University* 10:29–44.
- _____. 1957. A second look at the ants of the *Camponotus herculeanus* group in eastern Asia. *Journal of the Faculty of Agriculture, Kyushu University* 11:45–51.
- Yasumatsu, K., and Y. Murakami. 1960. A revision of the genus *Stenamma* of Japan (Hymenoptera, Formicidae, Myrmicinae). *Esakia* 1:27–31.
- Young, A. M. 1986. Notes on the distribution and abundance of ground- and arboreal-nesting ants (Hymenoptera: Formicidae) in some Costa Rican cacao habitats. *Proceedings of the Entomological Society of Washington* 88:550–571.
- Zhou, S.-Y., and Z.-M. Zheng. 1999. Taxonomic study of the ant genus *Pheidole* Westwood from Guangxi, with descriptions of three new species. *Acta Zootaxonomica Sinica* 24(1):83–88.
- Zimmerman, B. L., and M. T. Rodriguez. 1990. Frogs, snakes, and lizards of the INPA-WWF reserves near Manaus, Brazil. Pp. 426–454. In A. H. Gentry (ed.), *Four Neotropical Rainforests*. Yale University Press, New Haven, Connecticut.